

Anti-Skn1 antibody ab101724

[1 Abreviews](#) [画像数 1](#)

製品の概要

| | |
|--------------|---|
| 製品名 | Anti-Skn1 antibody |
| 製品の詳細 | Goat polyclonal to Skn1 |
| 由来種 | Goat |
| アプリケーション | 適用あり: WB |
| 種交差性 | 交差種: Human 交差が予測される動物種: Rat  |
| 免疫原 | Synthetic peptide corresponding to Human Skn1 aa 161-175 (internal sequence). Sequence: EPHLEASQHLPVPKH Database link: NP_055167.2  Run BLAST with  Run BLAST with |
| ポジティブ・コントロール | HEK293 and HepG2 cell lines. |
| 特記事項 | <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p> |

製品の特性

| | |
|-----------|---|
| 製品の状態 | Liquid |
| 保存方法 | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles. |
| バッファー | pH: 7.30 Preservative: 0.02% Sodium azide Constituents: 0.5% BSA, Tris buffered saline |
| 精製度 | Immunogen affinity purified |
| 特記事項 (精製) | Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. |

ポリモノ
アイソタイプ

ポリクローナル
IgG

アプリケーション


The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab101724の使用に適用されず
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

| アプリケーション | Abreviews | 特記事項 |
|----------|-----------|---|
| WB | | Use a concentration of 1 - 3 µg/ml. Predicted molecular weight: 47 kDa. 1 hour primary incubation is recommended for this product. |

ターゲット情報

| | |
|-------|--|
| 機能 | Transcription factor that binds to the octamer motif (5'-ATTTGCAT-3'). Regulated the expression of a number of genes such as SPRR2A or placental lactogen. |
| 組織特異性 | Specifically expressed in epidermis and cultured keratinocytes. |
| 配列類似性 | Belongs to the POU transcription factor family. Class-2 subfamily. Contains 1 homeobox DNA-binding domain. Contains 1 POU-specific domain. |
| 細胞内局在 | Nucleus. |

画像



Anti-Skn1 antibody (ab101724) at 2 µg/ml + HEK293 lysate in RIPA buffer at 35 µg

Predicted band size: 47 kDa

Incubation of ab101724 was 1 hour. Detected by chemiluminescence.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors