

Anti-SLC7A6OS antibody ab122727

1 References [画像数 3](#)

製品の概要

製品名	Anti-SLC7A6OS antibody
製品の詳細	Rabbit polyclonal to SLC7A6OS
由来種	Rabbit
アプリケーション	適用あり: IHC-P, WB, ICC/IF
種交差性	交差種: Human
免疫原	antigen sequence: AEALVLACKR LRSDAVESAA QKTSEGLERA AENNVFHLVA TVCSQEEPVQ PLLREVL RPS RDSQQRVRRN LRASAREVRQ EGRYRVLS, corresponding to amino acids 20-107 of Human SLC7A6OS. Run BLAST with ExpASY Run BLAST with NCBI
ポジティブ・コントロール	WB: Transfected HEK293T whole cells lysates; IHC-P: Human testis tissue; ICC: U-2 OS whole cells.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at -20°C or -80°C. Avoid freeze / thaw cycle. Store undiluted.
バッファー	pH: 7.20 Preservative: 0.02% Sodium azide Constituents: 40% Glycerol (glycerin, glycerine), 59% PBS
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab122727の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		1/500 - 1/1000. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
WB		Use a concentration of 0.04 - 0.4 µg/ml. Predicted molecular weight: 35 kDa.
ICC/IF		Use a concentration of 0.25 - 2 µg/ml. Fixation/Permeabilization: PFA/Triton X-100

ターゲット情報

配列類似性 Belongs to the SLC7A6OS family.

画像




Western blot - Anti-SLC7A6OS antibody (ab122727)

All lanes : Anti-SLC7A6OS antibody (ab122727) at 0.4 µg/ml

Lane 1 : Control (vector only transfected) - HEK-293T (human epithelial cell line from embryonic kidney transformed with large T antigen) whole cell lysate


Lane 2 : SLC7A6OS over-expression - HEK-293T whole cell lysate, Co-expressed with a C-terminal myc-DDK tag (~3.1 kDa)

Predicted band size: 35 kDa



ab122727, at 1/500 dilution, staining SLC7A6OS in paraffin-embedded Human testis tissue by Immunohistochemistry.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SLC7A6OS antibody (ab122727)



Immunocytochemistry/Immunofluorescence analysis of U-2 OS cells labelling SLC7A6OS with ab122727 shows positivity in the nucleus but is excluded from the nucleoli.

Immunocytochemistry/ Immunofluorescence - Anti-SLC7A6OS antibody (ab122727)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors