

Anti-SNAIL + SLUG antibody ab180714

★★★★★ [6 Abreviews](#) [185 References](#) [画像数 11](#)

製品の概要

製品名	Anti-SNAIL + SLUG antibody
製品の詳細	Rabbit polyclonal to SNAIL + SLUG
由来種	Rabbit
アプリケーション	適用あり: WB, IHC-P
種交差性	交差種: Mouse, Rat, Human, Recombinant fragment
免疫原	Synthetic peptide within Human SNAIL+SLUG aa 236-264. The exact sequence is proprietary. Database link: O95863
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at 4°C (stable for up to 12 months). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	pH: 7.30 Preservative: 0.02% Sodium azide Constituents: 50% Glycerol, 49% PBS
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab180714の使用に適用されます

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
WB	★★★★★ (2)	1/500 - 1/2000. Predicted molecular weight: 29 kDa.
IHC-P	★★★★★ (1)	1/50 - 1/200.

ターゲット情報


関連性

Function: SNAIL is involved in the epithelial to mesenchymal transition (EMT) and formation and maintenance of embryonic mesoderm (By similarity). Binds to 3 E-boxes of the E-cadherin gene promoter and represses its transcription. SLUG is a transcriptional repressor, involved in the generation and migration of neural crest cells. PTM: SNAIL is phosphorylated by GSK3B. Once phosphorylated, it becomes a target for BTRC ubiquitination. Ubiquitinated on Lys-98, Lys-137 and Lys-146 by FBXL14 and BTRC leading to degradation. BTRC-triggered ubiquitination requires previous GSK3B-mediated SNAI1 phosphorylation. Similarity: Both SNAIL and SLUG belong to the snail C2H2-type zinc-finger protein family. Tissue specificity: SNAIL is expressed in a variety of tissues with the highest expression in kidney. Expressed in mesenchymal and epithelial cell lines. SLUG is expressed in placenta and adult heart, pancreas, liver, kidney and skeletal muscle.

細胞内局在

Slug is generally nuclear, while Snail is known to be both cytoplasmic and nuclear. Once phosphorylated (probably on Ser-107, Ser-111, Ser-115 and Ser-119) snail is exported from the nucleus to the cytoplasm where subsequent phosphorylation of the destruction motif and ubiquitination involving BTRC occurs.

画像



Anti-SNAIL + SLUG antibody (ab180714) at 1/1000 dilution +
Mouse kidney at 25 µg


Secondary

HRP Goat Anti-Rabbit IgG (H+L) at 1/10000 dilution

Predicted band size: 29 kDa

Exposure time: 180 seconds

Blocking buffer: 3% nonfat dry milk in TBST.



Western blot - Anti-SNAIL + SLUG antibody
(ab180714)

All lanes : Anti-SNAIL + SLUG antibody (ab180714) at 1/1000 dilution

Lane 1 : A549 (Human lung carcinoma cell line) whole cell lysate

Lane 2 : 293T whole cell lysate

Lane 3 : HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate

Lysates/proteins at 25 µg per lane.


Secondary

All lanes : HRP Goat Anti-Rabbit IgG (H+L) at 1/10000 dilution

Predicted band size: 29 kDa

Exposure time: 90 seconds


Blocking buffer: 3% nonfat dry milk in TBST.



Western blot - Anti-SNAIL + SLUG antibody
(ab180714)

Anti-SNAIL + SLUG antibody (ab180714) at 1/500 dilution + Mouse heart tissue extracts

Predicted band size: 29 kDa



Western blot - Anti-SNAIL + SLUG antibody
(ab180714)

All lanes : Anti-SNAIL + SLUG antibody (ab180714)

Lane 1 : HeLa cell lysate


Lane 2 : A431 cell lysate

Lane 3 : HepG2 cell lysate

Lane 4 : Mouse lung tissue lysate


Lane 5 : Rat liver tissue lysate

Predicted band size: 29 kDa




Immunohistochemical analysis of paraffin-embedded mouse testis tissue labeling SNAIL1+SLUG with ab180714 at 1/50 dilution.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SNAIL + SLUG antibody
(ab180714)



Immunohistochemical analysis of paraffin-embedded rat brain tissue labeling SNAIL + SLUG with ab180714 at 1/100 dilution.


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SNAIL + SLUG antibody (ab180714)



ab180714 staining SNAIL in Mouse kidney tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde; antigen retrieval was by heat mediation in a basic buffer (TRIS+EDTA, pH9). Samples were incubated with primary antibody (1/500 in antibody diluent) for 1 hour. An undiluted HRP-conjugated Goat anti-rabbit IgG polyclonal was used as the secondary antibody.


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SNAIL + SLUG antibody (ab180714)

Image is courtesy of an anonymous Abreview




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SNAIL + SLUG antibody (ab180714)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human kidney tissue labelling SNAIL with ab180714. Tissue was fixed with formaldehyde and blocked with 5% BSA for 20 min at room temperature; antigen retrieval was by heat mediation with EDTA buffer (pH 9.0). Samples were incubated with primary antibody (1/100) for 12 hours at 4°C. A Biotin-conjugated mouse anti-rabbit IgG polyclonal (1/200) was used as the secondary antibody. Magnification: 200X.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SNAIL + SLUG antibody (ab180714)

Immunohistochemical analysis of paraffin-embedded human colon carcinoma tissue labeling SNAIL +SLUG with ab180714 at 1/100 dilution.



Western blot - Anti-SNAIL + SLUG antibody
(ab180714)


All lanes : Anti-SNAIL + SLUG antibody (ab180714)

Lane 1 : Snail recombinant with GST tag (55 kDa)

Lane 2 : Slug recombinant with GST tag (56 kDa)

Lysates/proteins at 0.1 µg per lane.

Predicted band size: 29 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-
embedded sections) - Anti-SNAIL + SLUG antibody
(ab180714)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human kidney tissue labelling SNAIL with ab180714. Tissue was fixed with formaldehyde and blocked with 5% BSA for 20 min at room temperature; antigen retrieval was by heat mediation with EDTA buffer (pH 9.0). Samples were incubated with primary antibody (1/100) for 12 hours at 4°C. A Biotin-conjugated mouse anti-rabbit IgG polyclonal (1/200) was used as the secondary antibody. Magnification: 400X.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors