


Anti-SNX16 antibody ab113885

画像数 4

製品の概要

製品名	Anti-SNX16 antibody
製品の詳細	Goat polyclonal to SNX16
由来種	Goat
アプリケーション	適用あり: WB, IHC-P
種交差性	交差種: Human 交差が予測される動物種: Mouse, Rat, Cow, Dog 
免疫原	Synthetic peptide (ATPYVPVPMPIGN-C) corresponding to N terminal amino acids 2-14 of Human SNX16. Run BLAST with Run BLAST with
ポジティブ・コントロール	Human breast, placenta and tonsil tissues; Daudi cell lysate
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.
バッファー	pH: 7.3 Preservative: 0.02% Sodium azide Constituents: 99% Tris buffered saline, 0.5% BSA
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab113885の使用に適用されず
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		Use a concentration of 0.05 - 0.2 µg/ml. Predicted molecular weight: 39 kDa.
IHC-P		Use a concentration of 3.75 µg/ml.

ターゲット情報

機能	May be involved in several stages of intracellular trafficking. Plays a role in protein transport from early to late endosomes. Plays a role in protein transport to the lysosome. Promotes degradation of EGFR after EGF signaling. Plays a role in intracellular transport of vesicular stomatitis virus nucleocapsids from the endosome to the cytoplasm.
組織特異性	Detected in placenta, lung, liver, heart and pancreas.
配列類似性	Belongs to the sorting nexin family. Contains 1 PX (phox homology) domain.
ドメイン	The PX domain mediates interaction with membranes enriched in phosphatidylinositol 3-phosphate.
細胞内局在	Early endosome membrane. Late endosome membrane. Cytoplasm. Lysosome.

画像




Anti-SNX16 antibody (ab113885) at 0.05 µg/ml + Daudi lysate (in RIPA buffer) at 35 µg

Developed using the ECL technique.


Predicted band size: 39 kDa

Exposure time: 1 hour




ab113885, at 3.75 $\mu\text{g/ml}$, staining SNX16 in formalin fixed, paraffin embedded Human breast tissue by Immunohistochemistry.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SNX16 antibody (ab113885)



ab113885, at 3.75 $\mu\text{g/ml}$, staining SNX16 in formalin fixed, paraffin embedded Human placenta tissue by Immunohistochemistry.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SNX16 antibody (ab113885)



ab113885, at 3.75 µg/ml, staining SNX16 in formalin fixed, paraffin embedded Human tonsil tissue by Immunohistochemistry.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SNX16 antibody (ab113885)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors