

# Anti-SNX30 antibody ab121600

**2 References** [画像数 3](#)

### 製品の概要

製品名	Anti-SNX30 antibody
製品の詳細	Rabbit polyclonal to SNX30
由来種	Rabbit
アプリケーション	<b>適用あり:</b> ICC/IF, IHC-P, WB
種交差性	<b>交差種:</b> Human
免疫原	antigen sequence: VGGDSTPSPD LLMARSGDK DLILPNGGTP AGTSSPASSS SLLNRLQLDD DIDGETRDLF VVDDPKKHV CTMETY, corresponding to amino acids 33-108 of Human SNX30. <a href="#">Run BLAST with</a> <a href="#">Expasy</a> <a href="#">Run BLAST with</a> <a href="#">NCBI</a>
ポジティブ・コントロール	Human duodenum tissue and RT4, U 251 MG, Human Liver and Tonsil lysates.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	pH: 7.20 Preservative: 0.02% Sodium azide Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine)
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

### アプリケーション

## The Abpromise guarantee

**Abpromise保証は、次のテスト済みアプリケーションにおけるab121600の使用に適用されます**

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF		Use a concentration of 0.25 - 2 µg/ml.
IHC-P		1/50 - 1/200. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
WB		Use a concentration of 0.04 - 0.4 µg/ml.

## ターゲット情報


### 機能

May be involved in several stages of intracellular trafficking.

### 配列類似性


Belongs to the sorting nexin family.  
Contains 1 PX (phox homology) domain.

## 画像




Immunofluorescent staining of Human cell line U-251MG shows positivity in nucleus, nucleoli and cytoplasm. Recommended concentration of ab121600 1-4 µg/ml. Cells treated with PFA/Triton X-100.

Immunocytochemistry/ Immunofluorescence - Anti-SNX30 antibody (ab121600)



ab121600 at 1/125 dilution staining SNX30 in Paraffin-embedded Human Duodenum tissue by Immunohistochemistry.  
Note: strong cytoplasmic positivity in glandular cells.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SNX30 antibody (ab121600)



**All lanes :** Anti-SNX30 antibody (ab121600) at 1/250 dilution

**Lane 1 :** RT4 cell lysate

**Lane 2 :** U 251 MG cell lysate

**Lane 3 :** Human Plasma

**Lane 4 :** Human Liver lysate

**Lane 5 :** Human Tonsil lysate

Developed using the ECL technique.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors