

Anti-Tenascin C antibody [EPR4219] - Low endotoxin, Azide free ab215369

リコンビナント RabMAb

3 References [画像数 7](#)

製品の概要

製品名	Anti-Tenascin C antibody [EPR4219] - Low endotoxin, Azide free
製品の詳細	Rabbit monoclonal [EPR4219] to Tenascin C - Low endotoxin, Azide free
由来種	Rabbit
特異性	IHC on human tissues which we tested (such as testis, pancreas and stomach) showed non-specific staining. We don't recommend this antibody for IHC on human tissues.
アプリケーション	適用あり: WB, IHC-P, IHC-Fr
種交差性	交差種: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	Human testis tissue; Human fetal brain lysate.
特記事項	<p>ab215369 is the carrier-free version of ab108930.</p> <p>Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit</p>

monoclonal antibodies. For details on our patents, please refer to [RabMAb® patents](#).

Our **Low endotoxin, azide-free formats** have low endotoxin level (≤ 1 EU/ml, determined by the LAL assay) and are free from azide, to achieve consistent experimental results in functional assays.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C. Do Not Freeze.
バッファー	pH: 7.20 Constituent: PBS
キャリア・フリー	はい
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR4219
アイソタイプ	IgG


アプリケーション

The **Abpromise guarantee** **Abpromise保証は、次のテスト済みアプリケーションにおけるab215369の使用に適用されず**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		Use at an assay dependent concentration. Predicted molecular weight: 241 kDa.
IHC-P		Use at an assay dependent concentration.
IHC-Fr		Use at an assay dependent concentration.

ターゲット情報

機能	Extracellular matrix protein implicated in guidance of migrating neurons as well as axons during development, synaptic plasticity as well as neuronal regeneration. Promotes neurite outgrowth from cortical neurons grown on a monolayer of astrocytes. Ligand for integrins alpha-8/beta-1, alpha-9/beta-1, alpha-V/beta-3 and alpha-V/beta-6.
配列類似性	Belongs to the tenascin family. Contains 15 EGF-like domains. Contains 1 fibrinogen C-terminal domain. Contains 15 fibronectin type-III domains.
細胞内局在	Secreted > extracellular space > extracellular matrix.




Immunohistochemistry (Frozen sections) - Anti-Tenascin C antibody [EPR4219] - Low endotoxin, Azide free (ab215369)

Immunohistochemistry (Frozen sections) analysis of rat cerebellar cortex labeling Tenascin C with **ab108930** at 1/100 dilution (4.27 µg/ml). Tissue was fixed with 4% PFA and permeabilized with 0.2% TritonX-100. Antigen retrieval was performed using a heated citrate solution (10mM citrate pH 6.0 + 0.05% Tween-20). **ab150077**, an Alexa Fluor® 488 Goat anti-Rabbit secondary antibody was used at 1/1000 (2 µg/ml). DAPI nuclear counterstain.

Positive staining on the molecular layer of rat cerebellar cortex (PMID: 1372043) is observed.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab108930**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Tenascin C antibody [EPR4219] - Low endotoxin, Azide free (ab215369)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse cerebellar cortex labeling Tenascin C with **ab108930** at 1/500 dilution (0.854 µg/ml). Heat mediated antigen retrieval was performed using Tris/EDTA buffer, pH 9 (**ab93684**). Hematoxylin was used to counterstain. A ready to use Goat Anti-Rabbit IgG H&L (HRP) secondary antibody was used. Staining on the molecular layer of mouse cerebellar cortex (PMID: 1372043) is observed.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab108930**).




Immunohistochemistry (Frozen sections) - Anti-Tenascin C antibody [EPR4219] - Low endotoxin, Azide free (ab215369)

Immunohistochemistry (Frozen sections) analysis of mouse E14 spinal cord labeling Tenascin C with **ab108930** at 1/100 dilution (4.27 µg/ml). Tissue was fixed with 4% PFA and permeabilized with 0.2% TritonX-100. Antigen retrieval was performed using a heated citrate solution (10mM citrate PH 6.0 + 0.05% Tween-20).

ab150077, an Alexa Fluor[®] 488 Goat anti-Rabbit secondary antibody was used at 1/1000 (2 µg/ml). DAPI nuclear counterstain.

Positive staining on mesenchymal condensations during chondrogenesis of mouse E14 embryo (PMID: 9822997; PMID: 19586317; PMID: 24778247) is observed.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab108930**).




Immunohistochemistry (Frozen sections) - Anti-Tenascin C antibody [EPR4219] - Low endotoxin, Azide free (ab215369)

Immunohistochemistry (Frozen sections) analysis of mouse E14 cerebellar cortex labeling Tenascin C with **ab108930** at 1/100 dilution (4.27 µg/ml). Tissue was fixed with 4% PFA and permeabilized with 0.2% TritonX-100. Antigen retrieval was performed using a heated citrate solution (10mM citrate pH 6.0 + 0.05% Tween-20). **ab150077**, an Alexa Fluor[®] 488 Goat anti-Rabbit secondary antibody was used at 1/1000 (2 µg/ml). DAPI nuclear counterstain.

Positive staining on the molecular layer of mouse E14 cerebellar cortex (PMID: 1372043) is observed.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab108930**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of rat cerebellar cortex labeling Tenascin C with **ab108930** at 1/500 dilution (0.854 µg/ml). Heat mediated antigen retrieval was performed using Tris/EDTA Buffer, pH 9 (**ab93684**). A ready to use Goat Anti-Rabbit IgG H&L (HRP) secondary antibody was used. Hematoxylin counterstain. Staining on the molecular layer of rat cerebellar cortex (PMID: 1372043) is observed.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab108930**).

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Tenascin C antibody [EPR4219] - Low endotoxin, Azide free (ab215369)



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse E14 spinal cord tissue sections labeling Tenascin C with **ab108930** at 1/500 dilution (0.854 µg/ml). Heat mediated antigen retrieval was performed using Tris/EDTA buffer, pH 9 (**ab93684**). A ready to use Goat Anti-Rabbit IgG H&L (HRP) secondary antibody was used. Hematoxylin counterstain.

Positive staining on mesenchymal condensations during chondrogenesis of mouse E14 embryo (PMID: 9822997; PMID: 19586317; PMID: 24778247) is observed.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab108930**).

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Tenascin C antibody [EPR4219] - Low endotoxin, Azide free (ab215369)

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Tenascin C antibody [EPR4219] - Low endotoxin, Azide free (ab215369)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors