


Anti-Tet2 antibody ab124297

KO 評価済

★★★★★ [10 Abreviews](#) [33 References](#) [画像数 6](#)

製品の概要

製品名	Anti-Tet2 antibody
製品の詳細	Rabbit polyclonal to Tet2
由来種	Rabbit
特異性	From Jan 2024, QC testing of replenishment batches of this polyclonal changed. All tested and expected application and reactive species combinations are still covered by our Abcam product promise. However, we no longer test all applications. For more information on a specific batch, please contact our Scientific Support who will be happy to help.
アプリケーション	適用あり: WB, ICC/IF
種交差性	交差種: Mouse 交差が予測される動物種: Human 
免疫原	Synthetic peptide corresponding to Mouse Tet2 aa 1600-1700 conjugated to keyhole limpet haemocyanin. Database link: Q4JK59
ポジティブ・コントロール	WB: E14Tg2A wild type mouse ES whole cell lysate.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.40 Preservative: 0.02% Sodium azide

Constituent: PBS

Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our scientific support team who will be happy to help.

精製度

Immunogen affinity purified

ポリ/モノ

ポリクローナル

アイソタイプ

IgG

アプリケーション

The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab124297の使用に適用されます

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB	★★★★★ (4)	1/250. Detects a band of approximately 223 kDa (predicted molecular weight: 223 kDa).
ICC/IF	★★★★★ (3)	Use a concentration of 1 µg/ml.

ターゲット情報

機能

Catalyzes the conversion of methylcytosine (5mC) to 5-hydroxymethylcytosine (hmC). Plays an important role in myelopoiesis. The clear function of 5-hydroxymethylcytosine (hmC) is still unclear but it may influence chromatin structure and recruit specific factors or may constitute an intermediate component in cytosine demethylation.

組織特異性

Broadly expressed. Highly expressed in hematopoietic cells; highest expression observed in granulocytes. Expression is reduced in granulocytes from peripheral blood of patients affected by myelodysplastic syndromes.

関連疾患

Note=TET2 is frequently mutated in myeloproliferative disorders (MPD). These constitute a heterogeneous group of disorders, also known as myeloproliferative diseases or myeloproliferative neoplasms (MPN), characterized by cellular proliferation of one or more hematologic cell lines in the peripheral blood, distinct from acute leukemia. Included diseases are: essential thrombocythemia, polycythemia vera, primary myelofibrosis (chronic idiopathic myelofibrosis). Bone marrow samples from patients display uniformly low levels of hmC in genomic DNA compared to bone marrow samples from healthy controls as well as hypomethylation relative to controls at the majority of differentially methylated CpG sites. Defects in TET2 are a cause of polycythemia vera (PV) [MIM:263300]. A myeloproliferative disorder characterized by abnormal proliferation of all hematopoietic bone marrow elements, erythroid hyperplasia, an absolute increase in total blood volume, but also by myeloid leukocytosis, thrombocytosis and splenomegaly.

Note=TET2 is frequently mutated in systemic mastocytosis; also known as systemic mast cell disease. A condition with features in common with myeloproliferative diseases. It is a clonal disorder of the mast cell and its precursor cells. The clinical symptoms and signs of systemic mastocytosis are due to accumulation of clonally derived mast cells in different tissues, including bone marrow, skin, the gastrointestinal tract, the liver, and the spleen.


Note=TET2 is frequently mutated in myelodysplastic syndromes, a heterogeneous group of

closely related clonal hematopoietic disorders. All are characterized by a hypercellular or hypocellular bone marrow with impaired morphology and maturation, dysplasia of the myeloid, megakaryocytic and/or erythroid lineages, and peripheral blood cytopenias resulting from ineffective blood cell production. Included diseases are: refractory anemia (RA), refractory anemia with ringed sideroblasts (RARS), refractory anemia with excess blasts (RAEB), refractory cytopenia with multilineage dysplasia and ringed sideroblasts (RCMD-RS). Chronic myelomonocytic leukemia (CMML) is a myelodysplastic/myeloproliferative disease. Myelodysplastic syndromes are considered a premalignant condition in a subgroup of patients that often progresses to acute myeloid leukemia (AML). Bone marrow samples from patients display uniformly low levels of hmC in genomic DNA compared to bone marrow samples from healthy controls as well as hypomethylation relative to controls at the majority of differentially methylated CpG sites.

配列類似性


Belongs to the TET family.

画像



ab124297 staining Tet2 in mES cells. The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.1% PBS-Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at 4°C with ab124297 at 1µg/ml and **ab7291**, Mouse monoclonal [DM1A] to alpha Tubulin - Loading Control. Cells were then incubated with **ab150081**, Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 488), pre-adsorbed at 1/1000 dilution (shown in green) and **ab150120**, Goat polyclonal Secondary Antibody to Mouse IgG - H&L (Alexa Fluor® 594), pre-adsorbed at 1/1000 dilution (shown in pseudocolour red). Nuclear DNA was labelled with DAPI (shown in blue).

Image was acquired with a confocal microscope (Leica-Microsystems TCS SP8) and a single confocal section is shown.



All lanes : Anti-Tet2 antibody (ab124297) at 1/250 dilution (Milk blocking - 1%)

Lane 1 : E14Tg2a (Mouse embryonic stem cell line) Whole Cell Lysate

Lane 2 : WT Mouse ES Cell Lysate (Positive Control for Tet2)

Lane 3 : Tet2 Knockout Mouse ES Cell Lysate (Negative Control)

Lysates/proteins at 25 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/10000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 223 kDa


Observed band size: 223 kDa

Additional bands at: 48 kDa (possible non-specific binding), 65 kDa (possible non-specific binding), 70 kDa (possible non-specific binding)

Exposure time: 4 minutes

This blot was produced using a 3-8% Tris Acetate gel under the TA buffer system. The gel was run at 150V for 60 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 1% Milk before being incubated with ab124297 overnight at 4°C. Antibody binding was detected using an anti-rabbit antibody conjugated to HRP, and visualised using ECL development solution.

Secondary antibody - goat **anti-rabbit HRP (ab97051)**



All lanes : Anti-Tet2 antibody (ab124297) at 1/1000 dilution
(Incubated for 12 hours at 4°C)

Lane 1 : Wild type mouse embryonic stem cells with Milk, 1 hour at 21°C

Lane 2 : Tet2 KO clone#1 mouse embryonic stem cells with Milk, 1 hour at 21°C

Lane 3 : Tet2 KO clone#2 mouse embryonic stem cells with Milk, 1 hour at 21°C

Lysates/proteins at 20 µg per lane.

Blocking peptides at 5 % per lane.

Secondary


All lanes : HRP conjugated Goat anti-rabbit IgG at 1/10000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 223 kDa


Exposure time: 5 minutes



Immunocytochemistry/ Immunofluorescence - Anti-Tet2 antibody (ab124297)

This image is courtesy of an anonymous abreview.

ICC/IF image of Anti-Tet2 antibody (ab124297) stained WT and Tet2^{-/-} mouse ES cells. The cells were fixed in PFA, permeabilized using 0.25% Triton X-100, and blocked with 10% BSA for 30 minutes. The cells were then incubated with ab124297 at a 1/400 dilution for 13 hours and 20 minutes at 4°C. The secondary antibody was a Rhodamine Red-X AffiniPure Donkey anti-Rabbit used at a 1/500 dilution.



Western blot - Anti-Tet2 antibody (ab124297)

This image is courtesy of an anonymous abreview.

All lanes : Anti-Tet2 antibody (ab124297) at 1/500 dilution

Lane 1 : Wild-type mouse embryonic stem cells

Lane 2 : Tet2^{-/-} mouse embryonic stem cells

Lysates/proteins at 20 μg per lane.

Secondary

All lanes : Goat anti-Rabbit (whole molecules) IgG HRP at 1/3000 dilution


Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 223 kDa

Observed band size: 250 kDa

Exposure time: 2 minutes



Immunocytochemistry/ Immunofluorescence - Anti-Tet2 antibody (ab124297)

This image is courtesy of an abreview from Joe Segal.

ICC/IF image of Anti-Tet2 antibody (ab124297) stained D3 mouse ES cells. The cells were fixed in 4% PFA, permeabilized using 0.1% Triton X-100, and blocked with 1% Goat serum, 0.1% BSA in PBS for 30 minutes. The cells were then incubated with ab124297 at a 1/100 dilution for 2 hours at RT. The secondary antibody was a Goat polyclonal Secondary Antibody to Rabbit IgG – H&L Alexa Fluor 488 (**ab150077**) used at a 1/200 dilution.

Secondary antibody - goat **anti-rabbit Alexa 488 (ab150077)**

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors