

# Anti-TEX12 antibody ab122455

**1 References** [画像数 3](#)

### 製品の概要

---

製品名	Anti-TEX12 antibody
製品の詳細	Rabbit polyclonal to TEX12
由来種	Rabbit
アプリケーション	<b>適用あり:</b> IHC-P, WB, ICC/IF
種交差性	<b>交差種:</b> Human
免疫原	antigen sequence: MANHLVKPDN RNCKRPRELE SPVPDSPQLS SLGKSDSSFS EISGLFYKDE ALEKDLNDVS, corresponding to amino acids 2-61 of Human TEX12.

 [Run BLAST with ExPASy](#)  [Run BLAST with NCBI](#)

**ポジティブ・コントロール** Human testis tissue.

**特記事項** The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

### 製品の特性

---

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	pH: 7.20 Preservative: 0.02% Sodium azide Constituents: 40% Glycerol (glycerin, glycerine), 59% PBS
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

### アプリケーション

---

## The Abpromise guarantee

**Abpromise保証は、次のテスト済みアプリケーションにおけるab122455の使用に適用されます**

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
IHC-P		1/200 - 1/500. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
WB		Use a concentration of 0.04 - 0.4 µg/ml.
ICC/IF		Use a concentration of 1 - 4 µg/ml. Recommend PFA Fixation and Triton X-100 treatment

## ターゲット情報

### 組織特異性


Testis specific.

## 画像



Immunofluorescent staining of Human cell line U-251 MG shows positivity in centrosome. Recommended concentration of ab122455 1-4 µg/ml. Cells treated with PFA/Triton X-100.

Immunocytochemistry/ Immunofluorescence - Anti-TEX12 antibody (ab122455)




Western blot - Anti-TEX12 antibody (ab122455)

**Lanes 2-3 :** Anti-TEX12 antibody (ab122455)

**Lane 1 :** Marker

**Lane 2 :** HEK293T cell lysate - vector only transfected

**Lane 3 :** HEK293T cell lysate - overexpressing TEX12 with a C-terminal myc-DDK tag (~3.1kDa)



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TEX12 antibody (ab122455)

ab122455, at 1/200 dilution, staining TEX12 in paraffin embedded Human testis tissue by Immunohistochemistry.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors