


Anti-TGF beta Receptor I antibody ab121024

★★★★★ [1 Abreviews](#) [7 References](#) [画像数 2](#)

製品の概要

製品名	Anti-TGF beta Receptor I antibody
製品の詳細	Goat polyclonal to TGF beta Receptor I
由来種	Goat
アプリケーション	適用あり: IHC-P, Flow Cyt (Intra), WB
種交差性	交差種: Rat, Human 交差が予測される動物種: Mouse, Rabbit, Horse, Cow, Dog, Pig, Rhesus monkey, Gorilla 
免疫原	Synthetic peptide within Human TGF beta Receptor I aa 50-150 (internal sequence). The exact immunogen sequence used to generate this antibody is proprietary information. If additional detail on the immunogen is needed to determine the suitability of the antibody for your needs, please contact our Scientific Support team to discuss your requirements. Corresponds to NP_004603.1 and NP_001124388.1. Database link: P36897 Run BLAST with Run BLAST with
ポジティブ・コントロール	WB: Rat liver and lung tissue lysates. IHC-P: Human adrenal gland and small intestine tissues. Flow Cyt (Intra): HeLa cells
特記事項	The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing. If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	pH: 7.30 Preservative: 0.02% Sodium azide Constituents: 99% Tris buffered saline, 0.5% BSA
精製度	Immunogen affinity purified

特記事項(精製)	ab121024 is purified from Goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
ポリモノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab121024の使用に適用されます
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		Use a concentration of 5 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
Flow Cyt (Intra)		Use a concentration of 10 µg/ml.
WB	★☆☆☆☆ (1)	Use a concentration of 0.03 - 0.1 µg/ml. Detects a band of approximately 55 kDa (predicted molecular weight: 55 kDa). 3% milk in TBST for blocking, 1 hr at RT. Primary antibody diluted in blocking buffer, 1 hr at room temperature.

ターゲット情報

機能	On ligand binding, forms a receptor complex consisting of two type II and two type I transmembrane serine/threonine kinases. Type II receptors phosphorylate and activate type I receptors which autophosphorylate, then bind and activate SMAD transcriptional regulators. Receptor for TGF-beta.
組織特異性	Found in all tissues examined, most abundant in placenta and least abundant in brain and heart.
関連疾患	Defects in TGFBR1 are the cause of Loeys-Dietz syndrome type 1A (LDS1A) [MIM:609192]; also known as Furlong syndrome or Loeys-Dietz aortic aneurysm syndrome (LDAS). LDS1 is an aortic aneurysm syndrome with widespread systemic involvement. The disorder is characterized by arterial tortuosity and aneurysms, craniosynostosis, hypertelorism, and bifid uvula or cleft palate. Other findings include exotropia, micrognathia and retrognathia, structural brain abnormalities, intellectual deficit, congenital heart disease, translucent skin, joint hyperlaxity and aneurysm with dissection throughout the arterial tree. Defects in TGFBR1 are the cause of Loeys-Dietz syndrome type 2A (LDS2A) [MIM:608967]. LDS2 is an aortic aneurysm syndrome with widespread systemic involvement. Physical findings include prominent joint laxity, easy bruising, wide and atrophic scars, velvety and translucent skin with easily visible veins, spontaneous rupture of the spleen or bowel, diffuse arterial aneurysms and dissections, and catastrophic complications of pregnancy, including rupture of the gravid uterus and the arteries, either during pregnancy or in the immediate postpartum period. LDS2 is characterized by the absence of craniofacial abnormalities with the exception of bifid uvula that can be present in some patients. Defects in TGFBR1 are the cause of aortic aneurysm familial thoracic type 5 (AAT5) [MIM:608967]. Aneurysms and dissections of the aorta usually result from degenerative changes in the aortic wall. Thoracic aortic aneurysms and dissections are primarily associated with a

characteristic histologic appearance known as 'medial necrosis' in which there is degeneration and fragmentation of elastic fibers, loss of smooth muscle cells, and an accumulation of basophilic ground substance.

配列類似性

Belongs to the protein kinase superfamily. TKL Ser/Thr protein kinase family. TGFB receptor subfamily.

Contains 1 GS domain.

Contains 1 protein kinase domain.


翻訳後修飾

Phosphorylated at basal levels in the absence of ligand binding. Activated by multiple phosphorylation, mainly in the GS region.

細胞内局在

Membrane.

画像



Lane 1 : Anti-TGF beta Receptor I antibody (ab121024) at 0.1 $\mu\text{g/ml}$

Lane 2 : Anti-TGF beta Receptor I antibody (ab121024) at 0.03 $\mu\text{g/ml}$

Lane 1 : Rat liver lysate in RIPA buffer


Lane 2 : Rat lung lysate in RIPA buffer

Lysates/proteins at 35 μg per lane.

Predicted band size: 55 kDa

Observed band size: 55 kDa

Primary incubation 1 hour at room temperature. Detected by chemiluminescence.



Flow Cytometry (Intracellular) - Anti-TGF beta
Receptor I antibody (ab121024)

Flow cytometric (Intracellular) analysis of paraformaldehyde fixed HeLa cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors