

Anti-Thyroglobulin antibody [1D4] ab16853

画像数 1

製品の概要

製品名	Anti-Thyroglobulin antibody [1D4]
製品の詳細	Mouse monoclonal [1D4] to Thyroglobulin
由来種	Mouse
特異性	Recognises thyroglobulin in hyperplastic and neoplastic thyroid.
アプリケーション	適用あり: IHC-P
種交差性	交差種: Human
免疫原	Full length protein corresponding to Human Thyroglobulin.
ポジティブ・コントロール	Thyroid
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.3 Preservative: 0.05% Sodium azide Constituents: Tissue culture supernatant, 1% BSA
精製度	Tissue culture supernatant
ポリ/モノ	モノクローナル
クローン名	1D4
アイソタイプ	IgG2a


The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab16853の使用に適用されず**
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		Use at an assay dependent concentration.

ターゲット情報

機能	Precursor of the iodinated thyroid hormones thyroxine (T4) and triiodothyronine (T3).
組織特異性	Thyroid gland specific.
関連疾患	Defects in TG are the cause of congenital hypothyroidism due to dysmorphogenesis type 3 (CHDH3) [MIM:274700]. A disorder due to thyroid dysmorphogenesis, causing large goiters of elastic and soft consistency in the majority of patients. Although the degree of thyroid dysfunction varies considerably among patients with defective thyroglobulin synthesis, patients usually have a relatively high serum free triiodothyronine (T3) concentration with disproportionately low free tetraiodothyronine (T4) level. The maintenance of relatively high free T3 levels prevents profound tissue hypothyroidism except in brain and pituitary, which are dependent on T4 supply, resulting in neurologic and intellectual defects in some cases. Variations in TG are associated with susceptibility to autoimmune thyroid disease type 3 (AITD3) [MIM:608175]. AITDs including Graves disease (GD) and Hashimoto thyroiditis (HT), are among the most common human autoimmune diseases. They are complex diseases, which are caused by an interaction between susceptibility genes and nongenetic factors, such as infection.
配列類似性	Belongs to the type-B carboxylesterase/lipase family. Contains 11 thyroglobulin type-1 domains.
翻訳後修飾	Sulfated tyrosines are desulfated during iodination.
細胞内局在	Secreted.

画像



Immunohistochemistry (Formalin fixed paraffin-embedded sections) analysis of human thyroid tissue labelling Amyloid A Component with ab16853.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Thyroglobulin antibody [1D4] (ab16853)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors