

Anti-TN-X antibody ab67586

★★★★★ [1 Abreviews](#) [画像数 1](#)

製品の概要

製品名	Anti-TN-X antibody
製品の詳細	Mouse polyclonal to TN-X
由来種	Mouse
アプリケーション	適用あり: WB
種交差性	交差種: Human
免疫原	Full length protein corresponding to Human TN-X aa 1-700. Database link: P22105 Run BLAST with Run BLAST with
ポジティブ・コントロール	TN-X transfected 293T cell lysate
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	pH: 7.40 Constituent: 100% PBS
精製度	Protein G purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab67586の使用に適用されます


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		1/500 - 1/1000. Detects a band of approximately 80 kDa (predicted molecular weight: 464 kDa).

ターゲット情報

機能	Appears to mediate interactions between cells and the extracellular matrix. Substrate-adhesion molecule that appears to inhibit cell migration. Accelerates collagen fibril formation. May play a role in supporting the growth of epithelial tumors.
組織特異性	Highly expressed in fetal adrenal, in fetal testis, fetal smooth, striated and cardiac muscle. Isoform XB-short is only expressed in the adrenal gland.
関連疾患	Tenascin-X deficiency (TNXD) [MIM:606408]: TNXD leads to an Ehlers-Danlos-like syndrome characterized by hyperextensible skin, hypermobile joints, and tissue fragility. Tenascin-X-deficient patients, however, lack atrophic scars, a major diagnostic criteria for classic Ehlers-Danlos. Delayed wound healing, which is also common in classic EDS, is only present in a subset of patients. Note=The disease is caused by mutations affecting the gene represented in this entry.
配列類似性	Belongs to the tenascin family. Contains 19 EGF-like domains. Contains 1 fibrinogen C-terminal domain. Contains 32 fibronectin type-III domains.
発生段階	Expression levels are lower in adults than in children.
細胞内局在	Secreted > extracellular space > extracellular matrix.

画像



All lanes : Anti-TN-X antibody (ab67586) at 1/500 dilution

Lane 1 : TN-X transfected 293T cell lysate

Lane 2 : Non-transfected 293T cell lysate

Lysates/proteins at 25 µg per lane.

Secondary

All lanes : Goat Anti-Mouse IgG (H&L)-HRP Conjugate at 1/2500 dilution

Predicted band size: 464 kDa

Observed band size: 80 kDa

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors