


# Anti-TNNI3K antibody ab111140

**2 References** [画像数 1](#)

### 製品の概要

製品名	Anti-TNNI3K antibody
製品の詳細	Rabbit polyclonal to TNNI3K
由来種	Rabbit
アプリケーション	<b>適用あり:</b> IHC-P
種交差性	<b>交差種:</b> Human <b>交差が予測される動物種:</b> Rat 
免疫原	Synthetic peptide derived from an internal region of Human TNNI3K.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
バッファー	<p>pH: 7.40</p> <p>Preservative: 0.02% Sodium azide</p> <p>Constituents: 49.1% PBS, 0.88% Sodium chloride, 50% Glycerol (glycerin, glycerine)</p> <p>PBS is without Mg<sup>2+</sup> and Ca<sup>2+</sup>.</p>
精製度	Immunogen affinity purified
特記事項 (精製)	ab111140 was affinity purified from rabbit antiserum by affinity chromatography using epitope specific immunogen.
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

## アプリケーション


**The Abpromise guarantee** Abpromise保証は、 次のテスト済みアプリケーションにおけるab111140の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		1/50 - 1/100. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

## ターゲット情報

機能	May play a role in cardiac physiology.
組織特異性	Highly expressed in both adult and fetal heart.
配列類似性	Belongs to the protein kinase superfamily. TKL Ser/Thr protein kinase family. MAP kinase kinase subfamily. Contains 10 ANK repeats. Contains 1 protein kinase domain.
翻訳後修飾	Autophosphorylated.
細胞内局在	Nucleus. Cytoplasm. Expressed at lower levels in the cytoplasm.

## 画像



ab111140, at a 1/50 dilution, staining TNNI3K in paraffin embedded Human lung carcinoma tissue by Immunohistochemistry. Image on the right was from tissue treated with the synthetic peptide.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TNNI3K antibody (ab111140)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors