


Anti-Transglutaminase 2 antibody [CUB 7402] ab2386

KO 評価済

★★★★★ [9 Abreviews](#) [34 References](#) [画像数 3](#)

製品の概要

製品名	Anti-Transglutaminase 2 antibody [CUB 7402]
製品の詳細	Mouse monoclonal [CUB 7402] to Transglutaminase 2
由来種	Mouse
アプリケーション	適用あり: IHC-P, WB, ICC/IF, Flow Cyt
種交差性	交差種: Mouse, Rabbit, Guinea pig, Cat, Dog, Human 交差が予測される動物種: Rat 
免疫原	Full length native protein (purified) corresponding to Transglutaminase 2. Purified from liver. Database link: P08587
エピトープ	Epitope mapping for this antibody is amino acid 447-478
ポジティブ・コントロール	IHC-P: Human breast carcinoma tissue. WB: A549 whole cell lysate; HeLa whole cell lysate; Huvec whole cell lysate
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.3 Preservative: 0.05% Sodium azide Constituent: 1% BSA
精製度	IgG fraction
ポリ/モノ	モノクローナル

クローン名	CUB 7402
ミエローマ	unknown
アイソタイプ	IgG1
軽鎖の種類	unknown

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab2386の使用に適用されます**


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
IHC-P	★★★★★ (3)	1/50 - 1/75. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
WB	★★★★★ (2)	Use a concentration of 1 µg/ml. Detects a band of approximately 77 kDa (predicted molecular weight: 77 kDa).
ICC/IF	★★★★★ (2)	Use at an assay dependent concentration. PubMed: 19398782
Flow Cyt	★★★★★ (1)	1/10. Methanol fixed cells. ab170190 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.

ターゲット情報

機能	Catalyzes the cross-linking of proteins and the conjugation of polyamines to proteins.
配列類似性	Belongs to the transglutaminase superfamily. Transglutaminase family.

画像



Western blot - Anti-Transglutaminase 2 antibody
[CUB 7402] (ab2386)

All lanes : Anti-Transglutaminase 2 antibody [CUB 7402] (ab2386)
at 1 µg/ml

Lane 1 : Wild-type A549 (Human lung carcinoma cell line) whole
cell lysate

Lane 2 : TGM2 knockout A549 (Human lung carcinoma cell line)
whole cell lysate

Lane 3 : HeLa (Human epithelial cell line from cervix
adenocarcinoma) whole cell lysate

Lane 4 : HUVEC (Human umbilical vein endothelial cell line) whole
cell lysate


Lysates/proteins at 20 µg per lane.

Predicted band size: 77 kDa

Observed band size: 77 kDa


Lanes 1 - 4: Merged signal (red and green). Green - ab2386
observed at 77 kDa. Red - loading control, **ab181602**, observed at
37 kDa.

ab2386 was shown to recognize in wild-type A549 cells as signal
was lost at the expected MW in TGM2 knockout cells. Additional
cross-reactive bands were observed in the wild-type and knockout
cells. Wild-type and TGM2 knockout samples were subjected to
SDS-PAGE. The membrane was blocked with 3% Milk. Ab2386
and **ab181602** (Rabbit anti-GAPDH loading control) were
incubated overnight at 4°C at 1 µg/ml and 1/20000 dilution
respectively. Blots were developed with Goat anti-Mouse IgG H&L
(IRDye® 800CW) preabsorbed **ab216772** and Goat anti-Rabbit IgG
H&L (IRDye® 680RD) preabsorbed **ab216777** secondary
antibodies at 1/20000 dilution for 1 hour at room temperature
before imaging.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Transglutaminase 2 antibody [CUB 7402] (ab2386)

Immunohistochemical analysis of human breast carcinoma tissue labeling Transglutaminase 2 with ab2386. Tissue was fixed with paraformaldehyde; antigen retrieval was by heat mediation in a citrate buffer. Samples were incubated with primary antibody (1/50) for 30 minutes at 22°C.



Flow Cytometry - Anti-Transglutaminase 2 antibody [CUB 7402] (ab2386)

Overlay histogram showing HCT116 cells stained with ab2386 (red line). The cells were fixed with methanol (5 min) and incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab2386, 1/10 dilution) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-mouse IgG (H+L) ([ab96879](#)) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG1 [ICIGG1] ([ab91353](#), 2µg/1x10⁶ cells) used under the same conditions. Acquisition of >5,000 events was performed. This antibody gave a decreased signal in HCT116 cells fixed with 4% paraformaldehyde (10 min) used under the same conditions.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors