

# Anti-TrkB antibody [EPR17805-146] ab187041

リコンビナント **RabMAb**

**1 Abreviews** **13 References** **画像数 9**

### 製品の概要

製品名	Anti-TrkB antibody [EPR17805-146]
製品の詳細	Rabbit monoclonal [EPR17805-146] to TrkB
由来種	Rabbit
アプリケーション	<b>適用あり:</b> Indirect ELISA, ICC/IF, WB, IHC-P, IP <b>適用なし:</b> Flow Cyt
種交差性	<b>交差種:</b> Mouse, Rat
免疫原	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: Mouse TrkB active protein; mouse and rat brain lysates. IHC-P: Mouse and rat cerebral cortex tissues and mouse primary neuron cell. IP: Mouse brain lysate. Indirect ELISA: Mouse TrkB antigen. ICC/IF: mouse primary neuron cells
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>.</p>

### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR17805-146

## アプリケーション

**The Abpromise guarantee** Abpromise保証は、次のテスト済みアプリケーションにおけるab187041の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Indirect ELISA		Use at an assay dependent concentration.
ICC/IF		1/50.
WB		1/5000. Detects a band of approximately 140, 90 kDa (predicted molecular weight: 92 kDa).
IHC-P		1/1000. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
IP		1/50.

## 追加情報

Is unsuitable for Flow Cyt.

## ターゲット情報

## 機能

Receptor for brain-derived neurotrophic factor (BDNF), neurotrophin-3 and neurotrophin-4/5 but not nerve growth factor (NGF). Involved in the development and/or maintenance of the nervous system. This is a tyrosine-protein kinase receptor. Known substrates for the TRK receptors are SHC1, PI-3 kinase, and PLC-gamma-1.

## 組織特異性

Isoform TrkB is widely expressed, mainly in the nervous tissue. In the CNS, expression is observed in the cerebral cortex, hippocampus, thalamus, choroid plexus, granular layer of the cerebellum, brain stem, and spinal cord. In the peripheral nervous system, it is expressed in many cranial ganglia, the opthalmic nerve, the vestibular system, multiple facial structures, the submaxillary glands, and dorsal root ganglia. Isoform TrkB-T1 is expressed in multiple tissues, mainly in brain, pancreas, kidney and heart. Isoform TrkB-T-Shc is predominantly expressed in brain.

## 配列類似性

Belongs to the protein kinase superfamily. Tyr protein kinase family. Insulin receptor subfamily. Contains 2 Ig-like C2-type (immunoglobulin-like) domains. Contains 2 LRR (leucine-rich) repeats. Contains 1 LRRCT domain. Contains 1 LRRNT domain. Contains 1 protein kinase domain.

## 翻訳後修飾


Ligand-mediated auto-phosphorylation.

## 細胞内局在

Membrane.


## 画像

### Indirect ELISA antibody dose-response curve antigen at 1000 ng/ml



Indirect ELISA - Anti-TrkB antibody [EPR17805-146] (ab187041)


ELISA analysis of Mouse TrkB (Ntrk2) recombinant protein at 1000 ng/ml with ab187041. An Alkaline Phosphatase-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L) at 1/2500 dilution was used as the secondary antibody.



Immunocytochemistry/ Immunofluorescence - Anti-TrkB antibody [EPR17805-146] (ab187041)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized mouse primary neuron cells labelling TrkB with ab187041 at 1/50 dilution, followed by **ab150077** Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) antibody at 1/1000 2 ug/ml dilution (Green). Confocal image showing positive staining in mouse primary neuron. Confocal scanning Z step was set as 0.3 µm followed by image processing with maximum Z projection. **ab11267** Anti-MAP2 mouse monoclonal antibody was used to counterstain tubulin at 1/500 4 dilution (Red). The Nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is **ab150077** Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 2 ug/ml dilution.



Western blot - Anti-TrkB antibody [EPR17805-146] (ab187041)

Anti-TrkB antibody [EPR17805-146] (ab187041) at 1/5000 dilution + Mouse TrkB active protein at 0.01 µg

### Secondary

Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/100000 dilution


**Predicted band size:** 92 kDa

**Observed band size:** 75-85 kDa

**Exposure time:** 3 seconds

Blocking/Dilution buffer: 5% NFDM/TBST.

Mouse TrkB active protein contains aa1-429 with a C-terminal poly histidine-tag.



**All lanes :** Anti-TrkB antibody [EPR17805-146] (ab187041) at 1/5000 dilution

**Lane 1 :** Mouse brain lysate

**Lane 2 :** Rat brain lysate

Lysates/proteins at 10 µg per lane.

### Secondary

**All lanes :** Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution

**Predicted band size:** 92 kDa


**Observed band size:** 140,90 kDa

**Exposure time:** 3 minutes

Blocking/Dilution buffer: 5% NFDm/TBST.

TrkB is abundantly expressed in the central and peripheral nervous system. The 140 kDa observed MW which is higher than the predicted one is due to the glycosylation modification.

90 kDa corresponds to the truncated form, as observed in the literature (PMID:24860020).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TrkB antibody [EPR17805-146] (ab187041)


Immunohistochemical analysis of paraffin-embedded mouse cerebral cortex tissue labeling TrkB with ab187041 at 1/1000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

Cytoplasmic and membrane staining on mouse cerebral cortex is observed.

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TrkB antibody [EPR17805-146] (ab187041)


Immunohistochemical analysis of paraffin-embedded rat cerebral cortex tissue labeling TrkB with ab187041 at 1/1000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

Cytoplasmic and membrane staining on rat cerebral cortex is observed.

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TrkB antibody  
[EPR17805-146] (ab187041)

Immunohistochemical analysis of paraffin-embedded mouse liver tissue labeling TrkB with ab187041 at 1/1000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

**Negative control:** no staining on mouse liver [PMID: 2555172].

Counter stained with Hematoxylin.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunoprecipitation - Anti-TrkB antibody  
[EPR17805-146] (ab187041)

TrkB was immunoprecipitated from 1 mg of mouse brain lysate with ab187041 at 1/50 dilution.

Western blot was performed from the immunoprecipitate using ab187041 at 1/1000 dilution.

VeriBlot for IP Detection Reagent (HRP) ([ab131366](#)), was used for detection at 1/10000 dilution.

Lane 1: Mouse brain lysate 10µg (Input).


Lane 2: ab187041 IP in mouse brain lysate.

Lane 3: Rabbit IgG, monoclonal [EPR25A] - Isotype Control ([ab172730](#)) instead of ab187041 in mouse brain lysate.

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 1 second.

### Why choose a recombinant antibody?



Anti-TrkB antibody [EPR17805-146] (ab187041)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors