


Anti-TRRAP antibody ab72509

画像数 1

製品の概要

製品名	Anti-TRRAP antibody
製品の詳細	Rabbit polyclonal to TRRAP
由来種	Rabbit
アプリケーション	適用あり: IP
種交差性	<p>交差種: Human</p> <p>交差が予測される動物種: Mouse, Rabbit, Chicken, Guinea pig, Cow, Dog, Pig, Chimpanzee, Rhesus monkey, Orangutan, Medaka fish, Platypus </p>
免疫原	<p>Synthetic peptide within Human TRRAP aa 3250-3350. The exact immunogen sequence used to generate this antibody is proprietary information. If additional detail on the immunogen is needed to determine the suitability of the antibody for your needs, please contact our Scientific Support team to discuss your requirements. AAD04629.1</p> <p>Database link: Q9Y4A5</p>
ポジティブ・コントロール	Whole cell lysate from Hela cells.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	<p>pH: 7</p> <p>Preservative: 0.09% Sodium azide</p> <p>Constituents: 1.815% Tris, 1.764% Sodium citrate, 0.021% PBS</p>
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab72509の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。


アプリケーション	Abreviews	特記事項
IP		Use at 2-5 µg/mg of lysate.

ターゲット情報

関連性 TRRAP (TRansformation/tRanscription domain Associated Protein) is an adapter protein that is found in a number of multiprotein chromatin complexes with histone acetyltransferase (HAT) activity. These complexes include PCAF, TFTC-HAT, TIP60 HAT, STAGA, and BAF53 complexes. TRRAP is thought to be responsible for the concerted and context-dependent recruitment of these complexes to chromatin during transcription, replication and DNA repair. TRRAP plays a key role in transcriptional activation by c-Myc, p53/TP53, E2F1 and E2F4.

細胞内局在 Nuclear

画像



Immunoprecipitation/ Western Blot of TRRAP.

Lane 1: ab72509 at 3µg/mg whole cell lysate.

Lane 2: Control IgG.

Another antibody to TRRAP was used at 1µg/ml for WB.

Whole cell lysate from Hela cells at 1mg for IP, 20% of IP loaded.

Chemiluminescence with an exposure time of 30 seconds.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors