

Anti-Tuberin antibody [Y320] - BSA and Azide free ab178441

KO 評価済 リコンビナント RabMAb

6 References 画像数 7

製品の概要

製品名	Anti-Tuberin antibody [Y320] - BSA and Azide free
製品の詳細	Rabbit monoclonal [Y320] to Tuberin - BSA and Azide free
由来種	Rabbit
特異性	This antibody recognises Tuberous sclerosis complex-2 (TSC2 also known as Tuberin. It is predicted to detect isoform 2-4 based on sequence homology.
アプリケーション	適用あり: Flow Cyt (Intra), IHC-P, WB
種交差性	交差種: Mouse, Rat, Human
免疫原	Synthetic peptide within Human Tuberin aa 1750-1850 (C terminal). The exact sequence is proprietary.
ポジティブ・コントロール	WB: HAP1 and HeLa whole cell lysates. Rat and mouse brain tissue lysates; Flow Cyt (intra): Jurkat cells; IHC: Rat and mouse liver tissue. Human lung carcinoma tissue.
特記事項	<p>ab178441 is the carrier-free version of ab32554.</p> <p>Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit</p>

monoclonal antibodies. For details on our patents, please refer to [RabMAb® patents](#).

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C. Do Not Freeze.
バッファー	Constituent: PBS
キャリア・フリー	はい
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	Y320
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、**次のテスト済みアプリケーションにおけるab178441の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt (Intra)		Use at an assay dependent concentration. ab199376 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
WB		Use at an assay dependent concentration. Detects a band of approximately 220 kDa (predicted molecular weight: 200 kDa).

ターゲット情報

機能	In complex with TSC1, inhibits the nutrient-mediated or growth factor-stimulated phosphorylation of S6K1 and EIF4EBP1 by negatively regulating mTORC1 signaling. Acts as a GTPase-activating protein (GAP) for the small GTPase RHEB, a direct activator of the protein kinase activity of mTORC1. Implicated as a tumor suppressor. Involved in microtubule-mediated protein transport, but this seems to be due to unregulated mTOR signaling. Stimulates weakly the intrinsic GTPase activity of the Ras-related proteins RAP1A and RAB5 in vitro. Mutations in TSC2 lead to constitutive activation of RAP1A in tumors.
組織特異性	Liver, brain, heart, lymphocytes, fibroblasts, biliary epithelium, pancreas, skeletal muscle, kidney, lung and placenta.
関連疾患	Defects in TSC2 are the cause of tuberous sclerosis type 2 (TSC2) [MIM:613254]. TSC2 is an autosomal dominant multi-system disorder that affects especially the brain, kidneys, heart, and skin. It is characterized by hamartomas (benign overgrowths predominantly of a cell or tissue type

that occurs normally in the organ) and hamartias (developmental abnormalities of tissue combination). Clinical symptoms can range from benign hypopigmented macules of the skin to profound mental retardation with intractable seizures to premature death from a variety of disease-associated causes.

Defects in TSC2 are a cause of lymphangioleiomyomatosis (LAM) [MIM:606690]. LAM is a progressive and often fatal lung disease characterized by a diffuse proliferation of abnormal smooth muscle cells in the lungs. It affects almost exclusively young women and can occur as an isolated disorder or in association with tuberous sclerosis complex.

Contains 1 Rap-GAP domain.

Phosphorylation at Ser-1387, Ser-1418 or Ser-1420 does not affect interaction with TSC1.

Phosphorylation at Ser-939 and Thr-1462 by PKB/AKT1 is induced by growth factor stimulation.


Cytoplasm. Membrane. At steady state found in association with membranes.

配列類似性

翻訳後修飾

細胞内局在

画像



All lanes : Anti-Tuberin antibody [Y320] ([ab32554](#)) at 1/5000 dilution (purified)

Lane 1 : HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate boiled

Lane 2 : HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate unboiled

Lane 3 : Mouse brain lysate boiled

Lane 4 : Mouse brain lysate unboiled

Lane 5 : Rat brain lysate boiled

Lane 6 : Rat brain lysate unboiled

Lysates/proteins at 20 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Predicted band size: 200 kDa

Observed band size: 200 kDa

The unboiled lysate are recommended to be used in SDS-PAGE, in order to prevent membrane protein aggregation.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab32554](#)).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Tuberin antibody [Y320] - BSA and Azide free (ab178441)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of rat liver tissue sections labeling Tuberin with purified **ab32554** at 1/100 dilution (2.97 µg/mL). Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0). Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32554**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Tuberin antibody [Y320] - BSA and Azide free (ab178441)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse liver tissue sections labeling Tuberin with purified **ab32554** at 1/100 dilution (2.97 µg/mL). Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0). Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32554**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Tuberin antibody [Y320] - BSA and Azide free (ab178441)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human lung carcinoma tissue sections labeling Tuberin with purified **ab32554** at 1/100 dilution (2.97 µg/mL). Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0). Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32554**).



Flow Cytometry (Intracellular) - Anti-Tuberin antibody [Y320] - BSA and Azide free (ab178441)

Intracellular Flow Cytometry analysis of Jurkat (Human T cell leukemia T lymphocyte) cells labeling Tuberin with purified **ab32554** at 1/30 dilution (10 µg/mL) (Red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32554**).



Western blot - Anti-Tuberin antibody [Y320] - BSA and Azide free (ab178441)

This WB data was generated using the same anti-Tuberin antibody clone, Y320, in a different buffer formulation (cat# [ab32554](#) - unpurified).

Lane 1: Wild-type HAP1 whole cell lysate (20 µg)

Lane 2: Tuberin knockout HAP1 whole cell lysate (20 µg)

Lane 3: HeLa whole cell lysate (20 µg)

Lanes 1 - 4: Merged signal (red and green). Green - [ab32554](#) observed at 180 kDa. Red - loading control, [ab18058](#), observed at 130 kDa.

[ab32554](#) was shown to specifically react with tuberin in wild-type HAP1 cells. No band was observed when tuberin knockout samples were used. Wild-type and tuberin knockout samples were subjected to SDS-PAGE. Ab32554 and [ab18058](#) (Mouse anti Vinculin loading control) were incubated overnight at 4°C at 1/1000 dilution and 1/10000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed [ab216773](#) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed [ab216776](#) secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Tuberin antibody [Y320] - BSA and Azide free (ab178441)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors