abcam

Product datasheet

Anti-Ugt2b antibody ab113433

2 References 画像数1

製品の概要

製品名 Anti-Ugt2b antibody

製品の詳細 Goat polyclonal to Ugt2b

由来種 Goat

特異性 ab113433 is not expected to cross-react with other Ugt2 proteins.

アプリケーション **適用あり:** WB 種交差性 交差種: Rat

免疫原 Synthetic peptide:

KLDFLSALEEVIDN

by a Cysteine residue linker, corresponding to internal sequence amino acids 423-436 of Rat

Ugt2b (NP 113721.3).

Run BLAST with Run BLAST with

ポジティブ・コントロール

Rat Kidney lysate.

特記事項 The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

> Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態

保存方法 Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

バッファー pH: 7.30

Preservative: 0.02% Sodium azide

Constituents: 99% Tris buffered saline, 0.5% BSA

精製度 Immunogen affinity purified

特記事項(精製) ab113433 was purified from Goat serum by ammonium sulphate precipitation followed by antigen

affinity chromatography using the immunizing peptide.

アイソタイプ

lgG

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab113433の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		Use a concentration of 0.1 - 0.3 µg/ml. Detects a band of approximately 50 kDa (predicted molecular weight: 61 kDa). Primary incubation time: 1 hour.

ターゲット情報

関連性

Ugt2b is an enzyme involved in conjugating lipophilic aglycon substrates with glucuronic acid [RGD]. Ugt2b is of major importance in the conjugation and subsequent elimination of potentially toxic xenobiotics and endogenous compounds acting on various endogenous steroids, especially etiocholanolone and androsterone.

細胞内局在

 $\label{thm:membrane} \mbox{Microsome membrane; Single-pass membrane protein. Endoplasmic reticulum membrane;}$

Single-pass membrane protein.

画像



Anti-Ugt2b antibody (ab113433) at 0.1 μ g/ml + Rat Kidney lysate (in RIPA buffer) at 35 μ g

Developed using the ECL technique.

Predicted band size: 61 kDa

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors