abcam

Product datasheet

Anti-UNG antibody ab47680

1 References 画像数 3

製品の概要

製品名 Anti-UNG antibody

製品の詳細 Rabbit polyclonal to UNG

由来種 Rabbit

特異性 This antibody is specific to UNG1. It will not cross react with UNG2.

アプリケーション 適用あり: WB, ICC/IF, IHC-P

種交差性 交差種: Mouse, Human

免疫原 A 13 amino acid peptide from near the amino terminus of human UNG1

ポジティブ・コントロール C2C12 cell lysate

特記事項 The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C.

バッファー pH: 7.2

Preservative: 0.02% Sodium azide

Constituent: PBS

精製度 Protein A purified

ポリ/モノ ポリクローナル

アイソタイプ lgG

アプリケーション


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		Use a concentration of 1 - 2 µg/ml. Detects a band of approximately 37 kDa (predicted molecular weight: 35 kDa).
ICC/IF		Use a concentration of 20 µg/ml.
IHC-P		Use a concentration of 2 μg/ml.

ターゲット情報

機能	Excises uracil residues from the DNA which can arise as a result of misincorporation of dUMP residues by DNA polymerase or due to deamination of cytosine.	
組織特異性	lsoform 1 is widely expressed with the highest expression in skeletal muscle, heart and testicles. lsoform 2 has the highest expression levels in tissues containing proliferating cells.	
関連疾患	Defects in UNG are a cause of immunodeficiency with hyper-lgM type 5 syndrome (HIGM5) [MIM:608106]. Hyper-lgM syndrome is a condition characterized by normal or increased serum lgM concentrations associated with low or absent serum lgG, lgA, and lgE concentrations. HIGM5 is associated with profound impairment in immunoglobulin (lg) class-switch recombination (CSR) at a DNA precleavage step.	
配列類似性	Belongs to the uracil-DNA glycosylase family.	
翻訳後修飾	Isoform 1 is processed by cleavage of a transit peptide.	
細胞内局在	Mitochondrion and Nucleus.	

画像




Lane 1 : Anti-UNG antibody (ab47680) at $0.5 \mu g/ml$ **Lane 2 :** Anti-UNG antibody (ab47680) at $1 \mu g/ml$ **Lane 3 :** Anti-UNG antibody (ab47680) at $2 \mu g/ml$

All lanes: C2C12 lysate

Lysates/proteins at 15 µg per lane.

Predicted band size: 35 kDa **Observed band size:** 37 kDa

Anti rabbit IgG secondary antibodies can be used.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-UNG antibody (ab47680)

<u>**ab37170**</u> at 2 μ g/ml staining UNG in human heart tissue section by Immunohistochemistry (Formalin/ PFA fixed paraffin-embedded tissue sections).

Immunocytochemistry/ Immunofluorescence - Anti-UNG antibody (ab47680) Immunofluorescence of UNG1 in Human Heart cells using ab47680 at 20 ug/ml.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors