

Anti-UQCRC2 antibody [EPR13051] ab203832

リコンビナント **RabMAb**

★★★★★ **1 Abreviews** **16 References** 画像数 10

製品の概要

製品名	Anti-UQCRC2 antibody [EPR13051]
製品の詳細	Rabbit monoclonal [EPR13051] to UQCRC2
由来種	Rabbit
アプリケーション	適用あり: Flow Cyt (Intra), IHC-P, WB, ICC/IF, IP
種交差性	交差種: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: Human fetal brain, fetal heart and fetal kidney lysates; HepG2, HEK-293, Jurkat, C6 and PC-12 cell lysates; Mouse heart, rat brain and rat kidney lysates. IHC-P: Human liver tissue. ICC/IF: HepG2 and Jurkat cells. Flow Cyt (intra): Jurkat cells. IP: HepG2 and HEK-293 whole cell lysates.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	<p>pH: 7.2</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA</p>
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR13051

アイソタイプIgG

アプリケーション


The Abpromise guarantee Abpromise保証は、 次のテスト済みアプリケーションにおけるab203832の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
Flow Cyt (Intra)		1/150.
IHC-P		1/150. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
WB	★★★★★ (1)	1/2000. Detects a band of approximately 48 kDa (predicted molecular weight: 48 kDa).
ICC/IF		1/300.
IP		1/50.

ターゲット情報

機能	This is a component of the ubiquinol-cytochrome c reductase complex (complex III or cytochrome b-c1 complex), which is part of the mitochondrial respiratory chain. The core protein 2 is required for the assembly of the complex.
配列類似性	Belongs to the peptidase M16 family. UQCRC2/QCR2 subfamily.
細胞内局在	Mitochondrion inner membrane.

画像



All lanes : Anti-UQCRC2 antibody [EPR13051] (ab203832) at 1/2000 dilution

Lane 1 : Human fetal brain lysate

Lane 2 : HepG2 (Human liver hepatocellular carcinoma) cell lysate

Lane 3 : HEK-293 (Human epithelial cells from embryonic kidney) cell lysate

Lane 4 : Jurkat (Human T cell leukemia cells from peripheral blood) cell lysate

Lysates/proteins at 20 µg per lane.

Secondary
All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/50000


dilution

Predicted band size: 48 kDa

Observed band size: 48 kDa

Exposure time: 3 seconds

Blocking/Dilution buffer: 5% NFDM/TBST.



Western blot - Anti-UQCRC2 antibody [EPR13051]
(ab203832)

All lanes : Anti-UQCRC2 antibody [EPR13051] (ab203832) at 1/2000 dilution

Lane 1 : Human fetal heart lysate

Lane 2 : Human fetal kidney lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG at 1/10000 dilution

Predicted band size: 48 kDa

Observed band size: 48 kDa

Exposure time: 3 seconds

Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-UQCRC2 antibody [EPR13051] (ab203832) at 1/2000 dilution

Lane 1 : Mouse heart lysate

Lane 2 : Rat brain lysate

Lane 3 : Rat kidney lysate

Lane 4 : C6 (Rat glial tumor cells) cell lysate

Lane 5 : PC-12 (Rat adrenal gland pheochromocytoma) cell lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/50000 dilution

Predicted band size: 48 kDa

Observed band size: 48 kDa

Exposure time: 5 seconds

Blocking/Dilution buffer: 5% NFDM/TBST.




Immunohistochemical analysis of paraffin-embedded Human liver tissue labeling UQCRC2 with ab203832 at 1/150 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution. Cytoplasmic staining on Human liver tissue is observed. Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-UQCRC2 antibody [EPR13051] (ab203832)



Immunofluorescent analysis of 100% methanol-fixed, 0.1% Triton X-100 permeabilized HepG2 (Human liver hepatocellular carcinoma) cells labeling UQCRC2 with ab203832 at 1/300 dilution, followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (**ab150077**) secondary antibody at 1/1000 dilution (green). Confocal image showing cytoplasmic staining on HepG2 cell line. The nuclear counterstain is DAPI (blue). Tubulin is detected with **ab7291** (anti-Tubulin mouse mAb) at 1/1000 dilution and **ab150120** (AlexaFluor®594 Goat anti-Mouse secondary) at 1/1000 dilution (red).

The negative controls are as follows:-


-ve control 1: ab203832 at 1/300 dilution followed by **ab150120** (AlexaFluor®594 Goat anti-Mouse secondary) at 1/1000 dilution.
 -ve control 2: **ab7291** (anti-Tubulin mouse mAb) at 1/1000 dilution followed by **ab150077** (Alexa Fluor®488 Goat Anti-Rabbit IgG H&L) at 1/1000 dilution.



Immunofluorescent analysis of 100% methanol-fixed, 0.1% Triton X-100 permeabilized Jurkat (Human T cell leukemia cells from peripheral blood) cells labeling UQCRC2 with ab203832 at 1/300 dilution, followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (**ab150077**) secondary antibody at 1/1000 dilution (green). Confocal image showing cytoplasmic staining on Jurkat cell line. The nuclear counterstain is DAPI (blue). Tubulin is detected with **ab7291** (anti-Tubulin mouse mAb) at 1/1000 dilution and **ab150120** (AlexaFluor®594 Goat anti-Mouse secondary) at 1/1000 dilution (red).


The negative controls are as follows:-

-ve control 1: ab203832 at 1/300 dilution followed by **ab150120** (AlexaFluor®594 Goat anti-Mouse secondary) at 1/1000 dilution.
 -ve control 2: **ab7291** (anti-Tubulin mouse mAb) at 1/1000 dilution followed by **ab150077** (Alexa Fluor®488 Goat Anti-Rabbit IgG H&L) at 1/1000 dilution.



Flow Cytometry (Intracellular) - Anti-UQCRC2 antibody [EPR13051] (ab203832)

Intracellular flow cytometric analysis of 4% paraformaldehyde-fixed Jurkat (Human T cell leukemia cells from peripheral blood) cells labeling UQCRC2 with ab203832 at 1/150 dilution (red) compared with a rabbit monoclonal IgG isotype control (**ab172730**; black) and an unlabelled control (cells without incubation with primary antibody and secondary antibody; blue). Goat anti rabbit IgG (FITC) at 1/150 dilution was used as the secondary antibody.




Immunoprecipitation - Anti-UQCRC2 antibody [EPR13051] (ab203832)

UQCRC2 was immunoprecipitated from 1mg of HepG2 (Human liver hepatocellular carcinoma) whole cell lysate with ab203832 at 1/50 dilution. Western blot was performed from the immunoprecipitate using ab203832 at 1/1000 dilution. Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG, was used as secondary antibody at 1/1500.

Lane 1: HepG2 whole cell lysate 10ug (Input). Lane 2: ab203832 IP in HepG2 whole cell lysate. Lane 3: Rabbit monoclonal IgG (**ab172730**) instead of ab203832 in HepG2 whole cell lysate.

Blocking and dilution buffer and concentration: 5% NFDm/TBST. Exposure time: 1 second.



Immunoprecipitation - Anti-UQCRC2 antibody
[EPR13051] (ab203832)

UQCRC2 was immunoprecipitated from 1mg of HEK-293 (Human epithelial cells from embryonic kidney) whole cell lysate with ab203832 at 1/50 dilution. Western blot was performed from the immunoprecipitate using ab203832 at 1/1000 dilution. Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG, was used as secondary antibody at 1/1500.

Lane 1: HEK-293 whole cell lysate 10ug (Input). Lane 2: ab203832 IP in HEK-293 whole cell lysate. Lane 3: Rabbit monoclonal IgG (**ab172730**) instead of ab203832 in HEK-293 whole cell lysate.

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 1 second.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-UQCRC2 antibody [EPR13051] (ab203832)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors