

Anti-Urokinase antibody ab24121

★★★★★ **3 Abreviews** **15 References**

製品の概要

製品名	Anti-Urokinase antibody
製品の詳細	Rabbit polyclonal to Urokinase
由来種	Rabbit
特異性	This antibody binds to human HMW-scuPA (54 kDa), HMW-tsuPA (52 kDa) and LMW-scuPA (33 kDa). This antibody is highly specific for urokinase, no crossreactions were found with other serine proteases (tissue plasminogen activator, plasmin, trypsin, chymotrypsin, elastase, thrombin). Human whole plasma or plasma proteins do not interfere with its binding to urokinase.
アプリケーション	適用あり: IHC-Fr, ELISA, IHC-P, ICC/IF, Dot blot, RIA, WB
種交差性	交差種: Human
免疫原	Full length protein (Human). Extensively purified native HMW-tcuPA and LMW-scuPA (mixture 50/50), isolated from human urine.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	pH: 8.00 Constituents: 0.16% Sodium phosphate, 0.38% Sodium borate, 0.87% Sodium chloride, 1% Dextran, 1% Mannitol
精製度	Affinity purified
特記事項 (精製)	Pooled antisera are passed over DEAE-cellulose to produce IgG-enriched fraction, which is subjected to absorption with immobilized total human serum proteins to remove non-specific antibodies. The affinity purified antibody is obtained by binding to immobilized native human

urokinase, followed by elution with acidic buffer, neutralization, dialysis, dispensing and lyophilization.

ポリ/モノ

ポリクローナル

アイソタイプ

IgG

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab24121の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-Fr		Use at an assay dependent concentration.
ELISA		Use at an assay dependent concentration.
IHC-P	★★★★★ (1)	Use at an assay dependent concentration.
ICC/IF		Use at an assay dependent concentration.
Dot blot		Use at an assay dependent concentration.
RIA		Use at an assay dependent concentration.
WB	★★★★★ (2)	Use at an assay dependent concentration.

ターゲット情報

機能	Specifically cleave the zymogen plasminogen to form the active enzyme plasmin.
組織特異性	Expressed in the prostate gland and prostate cancers.
配列類似性	Belongs to the peptidase S1 family. Contains 1 EGF-like domain. Contains 1 kringle domain. Contains 1 peptidase S1 domain.
翻訳後修飾	Phosphorylation of Ser-158 and Ser-323 abolishes proadhesive ability but does not interfere with receptor binding.
細胞内局在	Secreted.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors