

# Anti-Urokinase antibody [H77A10] ab28230

**3 References** [画像数 1](#)

### 製品の概要

---

|          |   |
|----------|---|
| 製品名      | Anti-Urokinase antibody [H77A10]                                  |
| 製品の詳細    | Mouse monoclonal [H77A10] to Urokinase                            |
| 由来種      | Mouse   |
| アプリケーション | <b>適用あり:</b> WB   |
| 種交差性     | <b>交差種:</b> Mouse   |
| 免疫原      | Recombinant full length protein corresponding to Mouse Urokinase. |

### 特記事項

This antibody was produced in a knockout mouse.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

### 製品の特性

---

|        |   |
|--------|---|
| 製品の状態  | Liquid  |
| 保存方法   | Shipped at 4°C. Upon delivery aliquot. Avoid freeze / thaw cycle.       |
| バッファー  | pH: 6.60<br>Constituents: 0.82% Sodium phosphate, 0.87% Sodium chloride |
| 精製度    | Protein A purified  |
| 一次抗体備考 | This antibody was produced in a knockout mouse.                         |
| ポリ/モノ  | モノクローナル   |
| クローン名  | H77A10  |
| アイソタイプ | IgG   |

### アプリケーション

---

**The Abpromise guarantee**      **Abpromise保証は、次のテスト済みアプリケーションにおけるab28230の使用に適用されず**


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

| アプリケーション | Abreviews | 特記事項  |
|----------|-----------|---|
| WB       |           | Use a concentration of 1 - 3 µg/ml. Use under non reducing condition. Predicted molecular weight: 48 kDa. |

## ターゲット情報

|       |   |
|-------|---|
| 機能    | Specifically cleave the zymogen plasminogen to form the active enzyme plasmin.  |
| 組織特異性 | Expressed in the prostate gland and prostate cancers.   |
| 配列類似性 | Belongs to the peptidase S1 family.<br>Contains 1 EGF-like domain.<br>Contains 1 kringle domain.<br>Contains 1 peptidase S1 domain. |
| 翻訳後修飾 | Phosphorylation of Ser-158 and Ser-323 abolishes proadhesive ability but does not interfere with receptor binding.                  |
| 細胞内局在 | Secreted.   |

## 画像



**All lanes :** Anti-Urokinase antibody [H77A10] (ab28230) at 3 µg/ml

**Lane 1 :** Mouse Urokinase (0.5ug)

**Lane 2 :** MUPA (0.5ug) + CPAI (2.5ug)

**Lane 3 :** Prestained standard

Performed under non-reducing conditions.

**Predicted band size:** 48 kDa

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors