abcam

Product datasheet

Anti-USO1 antibody [EPR2140Y] ab68137

יעלאעבע RabMAb

1 References 画像数 2

製品の概要

製品名 Anti-USO1 antibody [EPR2140Y]

製品の詳細 Rabbit monoclonal [EPR2140Y] to USO1

由来種 Rabbit

アプリケーション **適用あり:** WB

適用なし: Flow Cyt,ICC/IF or IHC-P

種交差性 交差種: Human, African green monkey

交差が予測される動物種: Mouse, Rat 🔷

免疫原 Synthetic peptide within Human USO1 aa 1-100 (N terminal). The exact sequence is proprietary.

ポジティブ・コントロール WB: HeLa cell lysate; 293T cell lysate; HepG2 cell lysate; Cos cell lysate

特記事項 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.

バッファー pH: 7.2

Preservative: 0.01% Sodium azide

Constituents: 9% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA, 50% Tissue culture

supernatant

精製度 Protein A purified

ポリ/モノ モノクローナル クローン名 **EPR2140Y**

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab68137の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		1/1000 - 1/5000. Detects a band of approximately 108 kDa (predicted molecular weight: 108 kDa).


追加情報

Is unsuitable for Flow Cyt,ICC/IF or IHC-P.

ターゲット情報

機能	General vesicular transport factor required for intercisternal transport in the Golgi stack; it is required for transcytotic fusion and/or subsequent binding of the vesicles to the target membrane. May well act as a vesicular anchor by interacting with the target membrane and holding the vesicular and target membranes in proximity.
配列類似性	Belongs to the VDP/USO1/EDE1 family. Contains 10 ARM repeats.
ドメイン	Composed of a globular head, an elongated tail (coiled-coil) and a highly acidic C-terminal domain.
翻訳後修飾	Phosphorylated in a cell cycle-specific manner; phosphorylated in interphase but not in mitotic cells. Dephosphorylated protein associates with the Golgi membrane; phosphorylation promotes dissociation.
細胞内局在	Cytoplasm > cytosol. Golgi apparatus membrane. Recycles between the cytosol and the Golgi apparatus during interphase. During interphase, the phosphorylated form is found exclusively in cytosol; the unphosphorylated form is associated with Golgi apparatus membranes.

画像



Western blot - Anti-USO1 antibody [EPR2140Y] (ab68137)

All lanes: Anti-USO1 antibody [EPR2140Y] (ab68137) at 1/500

dilution


Lane 1: HeLa cell lysate
Lane 2: 293T cell lysate
Lane 3: HepG2 cell lysate
Lane 4: Cos cell lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : HRP-labeled goat anti-rabbit at 1/2000 dilution

Predicted band size: 108 kDa **Observed band size:** 108 kDa



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors