

Anti-VAP1 antibody [174-5] ab81718

2 References [画像数 1](#)

製品の概要

製品名	Anti-VAP1 antibody [174-5]
製品の詳細	Mouse monoclonal [174-5] to VAP1
由来種	Mouse
アプリケーション	適用あり: IHC-Fr
種交差性	交差種: Human
免疫原	Tissue, cells or virus corresponding to Human VAP1.
ポジティブ・コントロール	Human tonsil tissue.
特記事項	<p>This antibody inhibits lymphocyte infiltration in liver allograft rejection in rats.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	Constituents: 0.1% BSA, PBS
特記事項 (精製)	ab81718 is purified and 0.2µM filtered.
一次抗体 備考	This antibody inhibits lymphocyte infiltration in liver allograft rejection in rats.
ポリ/モノ	モノクローナル
クローン名	174-5
アイソタイプ	IgG1

アプリケーション


The Abpromise guarantee **Abpromise保証は、**次のテスト済みアプリケーションにおけるab81718の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-Fr		1/50.

ターゲット情報

機能	Cell adhesion protein that participates in lymphocyte recirculation by mediating the binding of lymphocytes to peripheral lymph node vascular endothelial cells in an L-selectin-independent fashion. Has a monoamine oxidase activity. May play a role in adipogenesis.
組織特異性	Strongly expressed on the high endothelial venules of peripheral lymph nodes and on hepatic endothelia. Also highly expressed in appendix, lung and small intestine. Expressed also in adipose tissue, in bone marrow, colon, heart, kidney, ovary, pancreas, placenta, prostate, skeletal muscle, spleen and testis.
配列類似性	Belongs to the copper/topaquinone oxidase family.
翻訳後修飾	Topaquinone (TPQ) is generated by copper-dependent autoxidation of a specific tyrosyl residue. N- and O-glycosylated.
細胞内局在	Membrane.

画像



ab81718 at 1/50 dilution staining VAP1 in human tonsil tissue section by Immunohistochemistry (frozen sections).

Immunohistochemistry (Frozen sections) - Anti-VAP1 antibody [174-5] (ab81718)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors