# abcam

## Product datasheet

## Anti-Vimentin antibody ab24525

★★★★★ 11 Abreviews 99 References 画像数 4

#### 製品の概要

製品名 Anti-Vimentin antibody

製品の詳細 Chicken polyclonal to Vimentin

由来種 Chicken

アプリケーション 適用あり: ICC/IF, WB

種交差性 交差種: Mouse, Rat, Human

免疫原 Recombinant full length protein corresponding to Human Vimentin.

Database link: P08670

特記事項

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

#### 製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C. Do Not Freeze.

**バッファー** pH: 7.40

Preservative: 0.03% Sodium azide

Constituent: 99% PBS

精製度 lgY fraction ポリ/モノ ポリクローナル

アイソタイプ lgY

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab24525の使用に適用されます

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

1

アプリケーション	Abreviews	特記事項
ICC/IF	**** <u>(1)</u>	Use at an assay dependent concentration.
WB	*** <u>*</u>	1/10000.

## ターゲット情報

機能 Vimentins are class-Ill intermediate filaments found in various non-epithelial cells, especially

mesenchymal cells. Vimentin is attached to the nucleus, endoplasmic reticulum, and

mitochondria, either laterally or terminally.

Involved with LARP6 in the stabilization of type I collagen mRNAs for CO1A1 and CO1A2.

組織特異性 Highly expressed in fibroblasts, some expression in T- and B-lymphocytes, and little or no

expression in Burkitt's lymphoma cell lines. Expressed in many hormone-independent mammary

carcinoma cell lines.

**関連疾患** Cataract 30

配列類似性 Belongs to the intermediate filament family.

ドメイン The central alpha-helical coiled-coil rod region mediates elementary homodimerization.

The [IL]-x-C-x-x-[DE] motif is a proposed target motif for cysteine S-nitrosylation mediated by the

iNOS-S100A8/A9 transnitrosylase complex.

翻訳後修飾 Filament disassembly during mitosis is promoted by phosphorylation at Ser-55 as well as by

nestin (By similarity). One of the most prominent phosphoproteins in various cells of mesenchymal origin. Phosphorylation is enhanced during cell division, at which time vimentin filaments are

significantly reorganized. Phosphorylation by PKN1 inhibits the formation of filaments.

Phosphorylated at Ser-56 by CDK5 during neutrophil secretion in the cytoplasm. Phosphorylated

by STK33.

O-glycosylated during cytokinesis at sites identical or close to phosphorylation sites, this

interferes with the phosphorylation status.


S-nitrosylation is induced by interferon-gamma and oxidatively-modified low-densitity lipoprotein

(LDL(ox)) possibly implicating the iNOS-S100A8/9 transnitrosylase complex.

細胞内局在 Cytoplasm.


製品の状態 Vimentin is found in connective tissue and in the cytoskeleton.

画像




Immunocytochemistry/ Immunofluorescence - Anti-Vimentin antibody (ab24525)

Immunocytochemistry/ Immunofluorescence analysis of neuron/glial cultures labeling Vimentin with ab24525 (green) and GFAP with ab7260 (red). Vimentin is the sole cytoplasmic intermediate filament subunit expressed in fibroblasts, microglial and endothelial cells. The flattened cells in the middle of the image which appear green are fibroblasts. Astrocytes may express primarily GFAP, or both GFAP and vimentin, and so appear red (GFAP only) or golden yellow (GFAP and Vimentin). In cells which express both GFAP and vimentin, the two proteins assemble to produce heteropolymer filaments.



Immunocytochemistry/ Immunofluorescence - Anti-Vimentin antibody (ab24525)

Rat cerebral cortex cultures stained with chicken antibody to vimentin <u>ab24525</u> (green) and rabbit antibody to GFAP (red). Note flattened fibroblastic cells are mostly green (i.e. vimentin positive, GFAP negative), while clearly astrocytic cells, express both vimentin and GFAP and therefore appear golden or orange. Certain other cells express predominantly GFAP and therefore appear red.



Western blot of Rat whole brain extract, HeLa, SH-SY5Y, HEK293, and NIH/3T3 cells probed with ab24525, showing a single strong band at  $\sim 50$  kDa.



ab24525 staining Vimentin in rat smooth muscle cells from mesenteric artery by Immunocytochemistry/ Immunofluorescence. Cells were fixed with 4% paraformaldehyde in physiological saline solution (PSS) 4 min at 4°C and permeabilized with 0.3% Triton x100 before blocking with 2% BSA was done for 30 minutes at 20°C. Samples were incubated with primary antibody (1/300: in PSS with 2%BSA and 0.3% Triton X-100) for 14 hours at 4°C. An Abcam's ab6875, goat anti-chicken IgY Texas Red was used as secondary antibody at 1/400 dilution.

Vimentin antibody (ab24525)

This image is a courtesy of Anonymous Abreview

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

• We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.co.jp/abpromise">https://www.abcam.co.jp/abpromise</a> or contact our technical team.

### Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors