


Anti-Vimentin antibody [VI-10] ab20346

KO 評価済

★★★★★ [21 Abreviews](#) [113 References](#) [画像数 9](#)

製品の概要

製品名	Anti-Vimentin antibody [VI-10]
製品の詳細	Mouse monoclonal [VI-10] to Vimentin
由来種	Mouse
特異性	The antibody VI-10 reacts with vimentin, a 57 kDa intermediate filament expressed in variety of mesenchymal and mesodermal cell types.
アプリケーション	適用あり: WB, IHC-P, Flow Cyt, ICC/IF
種交差性	交差種: Mouse, Rat, Chicken, Human 交差が予測される動物種: Pig 
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	ICC/IF: HAP1 cells; Mouse dissociated neural precursor cells; Chicken postnatal erythrocytes; RBL rat basophilioc cells. IHC-P: Human skin tissue. WB: HAP1 cell lysate. Flow Cyt: NIH3T3 cells.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	pH: 8.0 Preservative: 0.097% Sodium azide Constituent: PBS
精製度	Proprietary Purification

特記事項(精製)	Purified by precipitation and chromatography.
ポリ/モノ	モノクローナル
クローン名	VI-10
ミエローマ	unknown
アイソタイプ	IgM
軽鎖の種類	unknown

アプリケーション

The Abpromise guarantee Abpromise保証は、 次のテスト済みアプリケーションにおけるab20346の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
WB	★★★★★ (4)	Use at an assay dependent concentration. Predicted molecular weight: 54 kDa.
IHC-P	★★★★★ (4)	Use a concentration of 1 - 5 µg/ml.
Flow Cyt	★★★★★ (1)	Use 1µg for 10 ⁶ cells. <u>ab91545</u> - Mouse monoclonal IgM, is suitable for use as an isotype control with this antibody.
ICC/IF	★★★★★ (7)	Use a concentration of 1 µg/ml. Staining technique: (a) fix cells for 10 min in methanol at -20°C and for 6 min in acetone at -20°C; (b) fix cells directly in methanol for 10 min at -20°C or in acetone for 10 min at -20°C.

ターゲット情報

機能	Vimentins are class-III intermediate filaments found in various non-epithelial cells, especially mesenchymal cells. Vimentin is attached to the nucleus, endoplasmic reticulum, and mitochondria, either laterally or terminally. Involved with LARP6 in the stabilization of type I collagen mRNAs for CO1A1 and CO1A2.
組織特異性	Highly expressed in fibroblasts, some expression in T- and B-lymphocytes, and little or no expression in Burkitt's lymphoma cell lines. Expressed in many hormone-independent mammary carcinoma cell lines.
関連疾患	Cataract 30
配列類似性	Belongs to the intermediate filament family.
ドメイン	The central alpha-helical coiled-coil rod region mediates elementary homodimerization. The [IL]-x-C-x-x-[DE] motif is a proposed target motif for cysteine S-nitrosylation mediated by the iNOS-S100A8/A9 transnitrosylase complex.
翻訳後修飾	Filament disassembly during mitosis is promoted by phosphorylation at Ser-55 as well as by nestin (By similarity). One of the most prominent phosphoproteins in various cells of mesenchymal origin. Phosphorylation is enhanced during cell division, at which time vimentin filaments are significantly reorganized. Phosphorylation by PKN1 inhibits the formation of filaments.

Phosphorylated at Ser-56 by CDK5 during neutrophil secretion in the cytoplasm. Phosphorylated by STK33.

O-glycosylated during cytokinesis at sites identical or close to phosphorylation sites, this interferes with the phosphorylation status.

S-nitrosylation is induced by interferon-gamma and oxidatively-modified low-density lipoprotein (LDL(ox)) possibly implicating the iNOS-S100A8/9 transnitrosylase complex.


Cytoplasm.

細胞内局在

製品の状態

Vimentin is found in connective tissue and in the cytoskeleton.

画像



All lanes : Anti-Vimentin antibody [VI-10] (ab20346) at 1 µg/ml

Lane 1 : Wild-type HAP1 whole cell lysate


Lane 2 : VIM (Vimentin) knockout HAP1 whole cell lysate

Lysates/proteins at 20 µg per lane.

Predicted band size: 54 kDa


Lanes 1 - 2: Merged signal (red and green). Green - ab20346 observed at 57 kDa. Red - loading control, **ab181602**, observed at 37 kDa.

ab20346 was shown to specifically react with Vimentin in wild-type HAP1 cells as signal was lost in VIM (Vimentin) knockout cells. Wild-type and VIM (Vimentin) knockout samples were subjected to SDS-PAGE. Ab20346 and **ab181602** (Rabbit anti-GAPDH loading control) were incubated overnight at 4°C at 1 µg/ml and 1/20000 dilution respectively. Blots were developed with Goat anti-Mouse IgG H&L (IRDye® 800CW) preabsorbed **ab216772** and Goat anti-Rabbit IgG H&L (IRDye® 680RD) preabsorbed **ab216777** secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.




ab20346 staining Vimentin in wild-type HAP1 cells (top panel) and Vimentin knockout HAP1 cells (bottom panel). The cells were fixed with 100% methanol (5min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with ab20346 at 1µg/ml dilution and **ab202272** at 1/250 dilution (shown in pseudo-color red) overnight at +4°C. The cells were then incubated with **ab150117** (Goat Anti-Mouse IgG H&L (Alexa Fluor® 488)) at 1/1000 dilution for 1 hour. Nuclear DNA was labelled in blue with DAPI.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).




Immunohistochemical analysis of formalin-fixed, paraffin-embedded human skin tissue labeling Vimentin with ab20346 at 5 ug/ml.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Vimentin antibody [VI-10] (ab20346)



Immunocytochemistry/ Immunofluorescence - Anti-Vimentin antibody [VI-10] (ab20346)


Immunocytochemical analysis of chicken postnatal erythrocytes labeling Vimentin with ab20346 at 1 ug/ml (green). Cells were counterstained with anti-alpha-tubulin (red) and DAPI (blue).



Immunocytochemistry/ Immunofluorescence - Anti-Vimentin antibody [VI-10] (ab20346)


ab20346 at a 1/500 dilution staining mouse dissociated neural precursor cells by immunocytochemistry. The antibody was incubated with the cells for 1 hour and then detected using a goat **anti-mouse Alexa-Fluor 568** (red stain) **ab175473**>**ab175473**. The image also shows a blue nuclear counterstain. Confocal image is shown with 2x zoom detail in top right corner.

This image is courtesy of an Abreview submitted by **Randal Moldich** on **22 February 2006**.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Vimentin antibody [VI-10] (ab20346)


Human normal skin. Staining is localised to cytoplasm. Left panel: with primary antibody at 1 µg/mL. Right panel: isotype control. Sections were stained using an automated system DAKO Autostainer Plus, at room temperature: sections were rehydrated and antigen retrieved with the Dako 3 in 1 AR buffers citrate pH6.1 in a DAKO PT Link. Slides were peroxidase blocked in 3% H₂O₂ in methanol for 10 mins. They were then blocked with Dako Protein block for 10 minutes (containing casein 0.25% in PBS) then incubated with primary antibody for 20 min and detected with Dako envision flex amplification kit for mouse for 30 minutes. Colorimetric detection was completed with Diaminobenzidine for 5 minutes. Slides were counterstained with Haematoxylin and coverslipped under DePeX. Please note that for manual staining we recommend to optimize the primary antibody concentration and incubation time (overnight incubation), and amplification may be required.



Western blot - Anti-Vimentin antibody [VI-10] (ab20346)

Image from Gao MQ et al, J Cell Sci. 2010 Oct 15;123(Pt 20):3507-14. Epub 2010 Sep 14, Fig 3. DOI 10.1242/jcs.072900


ab20346 used at a 1/1000 dilution in Western Blot. 5-20 µg of total protein from each sample was loaded.



Flow Cytometry - Anti-Vimentin antibody [VI-10] (ab20346)

Overlay histogram showing NIH3T3 cells stained with **ab20346** (red line). The cells were fixed with methanol (5 min) and then permeabilized with 0.1% PBS-Triton for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab20346, 1 µg/1x10⁶ cells) for 30 min at 22°C. The secondary antibody used was a goat **anti-mouse DyLight® 488** (IgG; H+L) (**ab96879**) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgM [ICIGM] (**ab91545**, 2 µg/1x10⁶ cells) used under the same conditions. Acquisition of >5,000 events was performed. This antibody gave a positive signal in NIH3T3 cells fixed with 4% paraformaldehyde (10 min)/permeabilized in 0.1% PBS-Triton used under the same

conditions.



Immunofluorescence staining of RBL rat basophilioc cell line with anti-Vimentin (VI-10). Nuclei are stained with DAPI (blue).

Immunocytochemistry/ Immunofluorescence - Anti-Vimentin antibody [VI-10] (ab20346)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors