abcam

Product datasheet

Anti-VPS4B/MIG1 antibody ab102687

3 Rafarancas

面偽粉 1

医薬用外劇物

製品の概要

製品名 Anti-VPS4B/MIG1 antibody

製品の詳細 Rabbit polyclonal to VPS4B/MIG1

由来種 Rabbit

アプリケーション **適用あり:** WB 種交差性 交差種: Human

交差が予測される動物種: Mouse, Rat, Cow, Zebrafish 🔷

免疫原 Recombinant fragment corresponding to Human VPS4B/MIG1 aa 176-403.

ポジティブ・コントロール 293T whole cell lysate

特記事項 The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

> Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態

保存方法 Shipped at 4°C. Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.

バッファー pH: 7.00

Preservative: 0.01% Thimerosal (merthiolate)

Constituents: 1.21% Tris, 0.75% Glycine, 20% Glycerol (glycerin, glycerine)

精製度 Immunogen affinity purified

ポリ/モノ ポリクローナル

アイソタイプ lgG

アプリケーション

The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab102687の使用に適用されます

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		1/3000. Predicted molecular weight: 49 kDa.

ターゲット情報

機能

Involved in late steps of the endosomal multivesicular bodies (MVB) pathway. Recognizes membrane-associated ESCRT-III assemblies and catalyzes their disassembly, possibly in combination with membrane fission. Redistributes the ESCRT-III components to the cytoplasm for further rounds of MVB sorting. MVBs contain intraluminal vesicles (ILVs) that are generated by invagination and scission from the limiting membrane of the endosome and mostly are delivered to lysosomes enabling degradation of membrane proteins, such as stimulated growth factor receptors, lysosomal enzymes and lipids. In conjunction with the ESCRT machinery also appears to function in topologically equivalent membrane fission events, such as the terminal stages of cytokinesis and enveloped virus budding (HIV-1 and other lentiviruses).

組織特異性

Ubiquitously expressed.

配列類似性

Belongs to the AAA ATPase family.

Contains 1 MIT domain.

ドメイン

The MIT domain serves as an adapter for ESCRT-III proteins. It forms an asymmetric three-helix bundle that binds amphipathic MIM (MIT interacting motif) helices along the groove between MIT helices 2 and 3 present in a subset of ESCRT-III proteins thus establishing the canonical MIM-MIT interaction. In an extended conformation along the groove between helices 1 and 3, also binds to a type-2 MIT interacting motif (MIM2).

翻訳後修飾


Phosphorylated upon DNA damage, probably by ATM or ATR.

細胞内局在

Prevacuolar compartment membrane. Late endosome membrane. Membrane-associated in the prevacuolar endosomal compartment. Localized in HIV-1 particles purified from acutely infected

cells.

画像



Anti-VPS4B/MIG1 antibody (ab102687) at 1/3000 dilution + 293T whole cell lysate at 30 µg

Predicted band size: 49 kDa

10% SDS PAGE

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors