

# Anti-WSTF antibody ab70263

**1 References** [画像数 3](#)

### 製品の概要

---

製品名	Anti-WSTF antibody
製品の詳細	Rabbit polyclonal to WSTF
由来種	Rabbit
アプリケーション	<b>適用あり:</b> WB, IP, IHC-P
種交差性	<b>交差種:</b> Human <b>交差が予測される動物種:</b> Rat, Cow, Dog, Pig, Chimpanzee, Rhesus monkey, Opossum, Orangutan 
免疫原	A region between residues 1 and 50 of Human WSTF (SwissProt entry Q9UIG0)
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 製品の特性

---

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	pH: 6.8 Preservative: 0.09% Sodium azide Constituents: 1.815% Tris, 1.764% Sodium citrate, 0.021% PBS
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

### アプリケーション

---

The Abpromise guarantee

**Abpromise保証は、次のテスト済みアプリケーションにおけるab70263の使用に適用されます**


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
WB		1/1000 - 1/10000. Detects a band of approximately 200 kDa (predicted molecular weight: 171 kDa).
IP		Use at 2-5 µg/mg of lysate.
IHC-P		1/100 - 1/500. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

## ターゲット情報

<b>機能</b>	Atypical tyrosine-protein kinase that plays a central role in chromatin remodeling and acts as a transcription regulator. Involved in DNA damage response by phosphorylating 'Tyr-142' of histone H2AX (H2AXY142ph). H2AXY142ph plays a central role in DNA repair and acts as a mark that distinguishes between apoptotic and repair responses to genotoxic stress. Essential component of the WICH complex, a chromatin remodeling complex that mobilizes nucleosomes and reconfigures irregular chromatin to a regular nucleosomal array structure. The WICH complex regulates the transcription of various genes, has a role in RNA polymerase I and RNA polymerase III transcription, mediates the histone H2AX phosphorylation at 'Tyr-142', and is involved in the maintenance of chromatin structures during DNA replication processes. In the complex, it mediates the recruitment of the WICH complex to replication foci during DNA replication. Also involved in vitamin D-coupled transcription regulation via its association with the WINAC complex, a chromatin-remodeling complex recruited by vitamin D receptor (VDR), which is required for the ligand-bound VDR-mediated transrepression of the CYP27B1 gene. In the WINAC complex, plays an essential role by targeting the complex to acetylated histones, an essential step for VDR-promoter association.
<b>組織特異性</b>	Ubiquitously expressed with high levels of expression in heart, brain, placenta, skeletal muscle and ovary.
<b>関連疾患</b>	Note=BAZ1B is located in the Williams-Beuren syndrome (WBS) critical region. WBS results from a hemizygous deletion of several genes on chromosome 7q11.23, thought to arise as a consequence of unequal crossing over between highly homologous low-copy repeat sequences flanking the deleted region. Haploinsufficiency of BAZ1B may be the cause of certain cardiovascular and musculo-skeletal abnormalities observed in the disease.
<b>配列類似性</b>	Belongs to the WAL family. BAZ1B subfamily. Contains 1 bromo domain. Contains 1 DDT domain. Contains 1 PHD-type zinc finger. Contains 1 WAC domain.
<b>発生段階</b>	Expressed at equal levels in 19-23 weeks old fetal tissues.
<b>ドメイン</b>	The N-terminal part (1-345), including the WAC domain and the C motif, mediates the tyrosine-protein kinase activity. The bromo domain mediates the specific interaction with acetylated histones.
<b>翻訳後修飾</b>	Phosphorylated upon DNA damage, probably by ATM or ATR.
<b>細胞内局在</b>	Nucleus. Accumulates in pericentromeric heterochromatin during replication. Targeted to replication foci throughout S phase via its association with PCNA.

## 画像



Western blot - Anti-WSTF antibody (ab70263)

**All lanes :** Anti-WSTF antibody (ab70263) at 0.05  $\mu\text{g/ml}$

**Lane 1 :** Whole cell lysate from HeLa cells at 5  $\mu\text{g}$


**Lane 2 :** Whole cell lysate from HeLa cells at 15  $\mu\text{g}$

**Lane 3 :** Whole cell lysate from HeLa cells at 50  $\mu\text{g}$

**Predicted band size:** 171 kDa


**Observed band size:** 200 kDa

**Additional bands at:** 140 kDa, 190 kDa, 300 kDa. We are unsure as to the identity of these extra bands.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human placenta tissue labelling WSTF with ab70263 at 1/200 (1  $\mu\text{g/ml}$ ). Detection: DAB.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-WSTF antibody (ab70263)



Immunoprecipitation - Anti-WSTF antibody (ab70263)

1 mg of whole cell lysate from HeLa cells was immunoprecipitated using ab70263 at 3  $\mu\text{g/mg}$  lysate (lane 1) or a control rabbit Ig (lane 2). The subsequent blot was carried with ab70263 at 0.05  $\mu\text{g/ml}$ .

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors