


Anti-XBP1 antibody ab37152

★★★★★ [28 Abreviews](#) [146 References](#) [画像数 13](#)

製品の概要

製品名	Anti-XBP1 antibody
製品の詳細	Rabbit polyclonal to XBP1
由来種	Rabbit
特異性	XBP-1 antibody will recognize both isoforms (non-spliced and spliced) of XBP-1.
アプリケーション	適用あり: IHC-P, IHC-Fr, ICC, IHC (PFA fixed), WB, ICC/IF
種交差性	交差種: Mouse, Rat, Human 交差が予測される動物種: Cow 
免疫原	Synthetic peptide corresponding to Human XBP1 (N terminal). Synthetic peptide corresponding to 18 amino acids near to the N terminus of the human XBP1 protein. The immunogen is located within amino acids 40 - 90. Database link: P17861 (Peptide available as ab39910)
ポジティブ・コントロール	WB: HepG2 cell lysate, A549 cell lysate. IHC-P: Human liver and non-lesional skin tissue; Mouse testis and spleen tissue, rat liver tissue. ICC/IF: HepG2 cells.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	pH: 7.2 Preservative: 0.02% Sodium azide Constituent: PBS
精製度	Immunogen affinity purified

ポリモノ
アイソタイプ

ポリクローナル
IgG

アプリケーション


The Abpromise guarantee **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab37152の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P	★★★★★ (12)	Use a concentration of 5 µg/ml.
IHC-Fr		Use at an assay dependent concentration. PubMed: 20028872
ICC		Use a concentration of 10 µg/ml.
IHC (PFA fixed)		Use a concentration of 20 µg/ml.
WB	★★★★★ (11)	Use a concentration of 0.5 - 2 µg/ml. Predicted molecular weight: 29 kDa. Can be blocked with Human XBP1 peptide (ab39910) . Human samples only. Not batch tested in mouse or rat, please see AbReviews for more information. Use at a concentration of 1 - 2 µg/ml. Predicted molecular weight: 29 kDa (isoform 1) and 40 kD (isoform 2). Can be blocked with XBP1 peptide (ab39910).
ICC/IF	★★★★★ (3)	Use a concentration of 20 µg/ml.

ターゲット情報


機能	Transcription factor essential for hepatocyte growth, the differentiation of plasma cells, the immunoglobulin secretion, and the unfolded protein response (UPR). Acts during endoplasmic reticulum stress (ER) by activating unfolded protein response (UPR) target genes via direct binding to the UPR element (UPRE). Binds DNA preferably to the CRE-like element 5'-GATGACGTG[TG]N(3)[AT]T-3', and also to some TPA response elements (TRE). Binds to the HLA DR-alpha promoter. Binds to the Tax-responsive element (TRE) of HTLV-I.
関連疾患	Genetic variations in XBP1 could be associated with susceptibility to major affective disorder type 7 (MAFD7) [MIM:612371]. Major affective disorders represent a class of mental disorders characterized by a disturbance in mood as their predominant feature.
配列類似性	Belongs to the bZIP family. Contains 1 bZIP domain.
細胞内局在	Nucleus.

画像



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-XBP1 antibody (ab37152)

Immunohistochemical analysis of paraffin-embedded Rat Liver Tissue using ab37152 at 5 µg/ml. Tissue was fixed with formaldehyde and blocked with 10% serum for 1 h at RT; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody overnight at 4°C. A goat anti-rabbit IgG H&L (HRP) at 1/250 was used as secondary. Counter stained with Hematoxylin.



Western blot - Anti-XBP1 antibody (ab37152)

Lane 1 : Anti-XBP1 antibody (ab37152) at 0.25 µg/ml

Lane 2 : Anti-XBP1 antibody (ab37152) at 0.5 µg/ml

Lane 3 : Anti-XBP1 antibody (ab37152) at 1 µg/ml

All lanes : A549 cell lysate


Lysates/proteins at 15 µg per lane.

Secondary

All lanes : Goat anti-rabbit IgG HRP conjugate at 1/10000 dilution

Predicted band size: 29 kDa

1h incubation at RT in 5% NFDM/TBST.



Western blot - Anti-XBP1 antibody (ab37152)

Lane 1 : Anti-XBP1 antibody (ab37152) at 1 µg/ml

Lane 2 : Anti-XBP1 antibody (ab37152) at 2 µg/ml


All lanes : HepG2 cell lysate

Lysates/proteins at 15 µg per lane.

Secondary


All lanes : Goat anti-rabbit IgG HRP conjugate at 1/10000 dilution

Predicted band size: 29 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-XBP1 antibody (ab37152)

Immunohistochemistry of mouse testis cells labeling XBP1 with Anti-XBP1 antibody (ab37152) at 2µg/ml. Tissue was fixed with formaldehyde and blocked with 10% serum for 1 h at RT; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody overnight at 4°C. A goat anti-rabbit IgG H&L (HRP) at 1/250 was used as secondary. Counter stained with Hematoxylin.




Immunohistochemistry (PFA fixed) - Anti-XBP1 antibody (ab37152)

Immunohistochemistry of human liver tissue labeling XBP1 with Anti-XBP1 antibody (ab37152) at 20µg/ml. followed by goat anti-rabbit IgG secondary antibody at 1/500 dilution (green). DAPI staining (Blue).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-XBP1 antibody (ab37152)

Immunohistochemistry of mouse spleen cells labeling XBP1 with Anti-XBP1 antibody (ab37152) at 2µg/ml. Tissue was fixed with formaldehyde and blocked with 10% serum for 1 h at RT; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody overnight at 4°C. A goat anti-rabbit IgG H&L (HRP) at 1/250 was used as secondary. Counter stained with Hematoxylin.




Immunohistochemistry (PFA fixed) - Anti-XBP1 antibody (ab37152)

Immunohistochemistry of mouse pancreas tissue labeling XBP1 with Anti-XBP1 antibody (ab37152) at 20µg/ml. followed by goat anti-rabbit IgG secondary antibody at 1/500 dilution (green). DAPI staining (Blue).




Immunohistochemistry (PFA fixed) - Anti-XBP1 antibody (ab37152)

Immunohistochemistry of human pancreas tissue labeling XBP1 with Anti-XBP1 antibody (ab37152) at 20µg/ml. followed by goat anti-rabbit IgG secondary antibody at 1/500 dilution (green). DAPI staining (Blue).




Immunocytochemistry of HepG2 cells labeling XBP1 with Anti-XBP1 antibody (ab37152) at 20µg/ml. Blue: DAPI staining.

Immunocytochemistry/ Immunofluorescence - Anti-XBP1 antibody (ab37152)



Immunohistochemistry (Formalin-fixed paraffin embedded sections) of human liver tissue labeling XBP1 with Anti-XBP1 antibody (ab37152) at 5µg/ml.


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-XBP1 antibody (ab37152)



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-XBP1 antibody (ab37152)

Image courtesy of an anonymous Abreview.


ab37152 staining XBP1 in human liver tissue by Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections). Tissue was fixed with formaldehyde and a heat mediated antigen retrieval step was performed using Citrate-EDTA antigen retrieval buffer and microwave. Heating was performed in the microwave 3 minutes on high until boiling, 6 minutes on 20% power. Samples were then blocked for 15 minutes at 27°C followed by incubation with the primary antibody at a 1/100 dilution for 12 hours at 4°C. An undiluted HRP-conjugated rat polyclonal was used as the secondary antibody.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-XBP1 antibody (ab37152)

Image from Ren Yunqing et al., PLoS Genet. 2009 June; 5(6): e1000523 (Fig 6C).

ab37152 at 1/200 dilution staining XBP1 in human non-lesional skin tissue sections by Immunohistochemistry (Formalin/ PFA fixed paraformaldehyde tissue sections). A biotin labelled secondary was used at 1/100 dilution. Diaminobenzidine (DAB) was used for staining development in a ABC staining Kit and the sections were counterstained with haematoxylin.



Immunocytochemistry of XBP-1 in HepG2 cells with ab37152 at 10 ug/mL

Immunocytochemistry - Anti-XBP1 antibody
(ab37152)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors