




### Anti-YAP1 (phospho Y357) antibody ab62751

★★★★★ [1 Abreviews](#) [47 References](#) [画像数 1](#)

#### 製品の概要

製品名	Anti-YAP1 (phospho Y357) antibody
製品の詳細	Rabbit polyclonal to YAP1 (phospho Y357)
由来種	Rabbit
アプリケーション	<b>適用あり:</b> WB, ICC/IF
種交差性	<b>交差種:</b> Human <b>交差が予測される動物種:</b> Mouse, Rat, Horse, Chicken, Cow, Dog, Xenopus laevis, Chimpanzee, Zebrafish, Opossum 
免疫原	Synthetic peptide corresponding to Human YAP1 aa 350-450 (C terminal) (phospho Y357) conjugated to keyhole limpet haemocyanin. Database link: <a href="#">P46937</a> <div>  <a href="#">Run BLAST with</a>  <a href="#">Run BLAST with</a> </div>
ポジティブ・コントロール	HEK-293T cells co-transfected with human YAP1 and human c-Abl.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
バッファー	pH: 7.40 Preservative: 0.097% Sodium azide Constituent: 0.0268% PBS
精製度	Immunogen affinity purified
特記事項 (精製)	ab62751 was affinity-purified using the immunizing peptide immobilized on agarose.
ポリ/モノ	ポリクローナル

## アプリケーション


**The Abpromise guarantee** Abpromise保証は、次のテスト済みアプリケーションにおけるab62751の使用に適用されます  
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		Use a concentration of 0.5 - 1 µg/ml. Detects a band of approximately 65 kDa (predicted molecular weight: 65 kDa).
ICC/IF	★★★★★ (1)	Use at an assay dependent concentration.

## ターゲット情報

機能	Transcriptional regulator which can act both as a coactivator and a corepressor and is the critical downstream regulatory target in the Hippo signaling pathway that plays a pivotal role in organ size control and tumor suppression by restricting proliferation and promoting apoptosis. The core of this pathway is composed of a kinase cascade wherein MST1/MST2, in complex with its regulatory protein SAV1, phosphorylates and activates LATS1/2 in complex with its regulatory protein MOB1, which in turn phosphorylates and inactivates YAP1 oncoprotein and WWTR1/TAZ. Plays a key role to control cell proliferation in response to cell contact. Phosphorylation of YAP1 by LATS1/2 inhibits its translocation into the nucleus to regulate cellular genes important for cell proliferation, cell death, and cell migration. The presence of TEAD transcription factors are required for it to stimulate gene expression, cell growth, anchorage-independent growth, and epithelial mesenchymal transition (EMT) induction. Isoform 2 and isoform 3 can activate the C-terminal fragment (CTF) of ERBB4 (isoform 3).
組織特異性	Increased expression seen in some liver and prostate cancers. Isoforms lacking the transactivation domain found in striatal neurons of patients with Huntington disease (at protein level).
配列類似性	Belongs to the YORKIE family. Contains 2 WW domains.
翻訳後修飾	Phosphorylated by LATS1 and LATS2; leading to cytoplasmic translocation and inactivation. Phosphorylated by ABL1; leading to YAP1 stabilization, enhanced interaction with TP73 and recruitment onto proapoptotic genes; in response to DNA damage.
細胞内局在	Cytoplasm. Nucleus. Both phosphorylation and cell density can regulate its subcellular localization. Phosphorylation sequesters it in the cytoplasm by inhibiting its translocation into the nucleus. At low density, predominantly nuclear and is translocated to the cytoplasm at high density.

## 画像



**All lanes :** Anti-YAP1 (phospho Y357) antibody (ab62751) at 1/2000 dilution

**Lane 1 :** Whole cell lysates of HEK-293T cells co-transfected with human YAP1 and human c-Abl

**Lane 2 :** Whole cell lysates of untransfected HEK-293T cells

**Lane 3 :** Whole cell lysates of HEK-293T cells co-transfected with human YAP1 and human c-Abl with YAP1 peptide (human 351-362) at 20 µg/ml

**Lane 4 :** Whole cell lysates of HEK-293T cells co-transfected with human YAP1 and human c-Abl with phospho-YAP1 (human 351-362 [phospho Y357]) immunizing peptide at 20 µg/ml

### Secondary

**All lanes :** Goat Anti-Rabbit IgG, Peroxidase conjugate

**Predicted band size:** 65 kDa

**Observed band size:** 65 kDa

**Additional bands at:** 100 kDa, 70 kDa. We are unsure as to the identity of these extra bands.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors