

Anti-ZNF787 antibody ab117837

画像数 2

製品の概要

製品名	Anti-ZNF787 antibody
製品の詳細	Rabbit polyclonal to ZNF787
由来種	Rabbit
アプリケーション	適用あり: WB, IP
種交差性	交差種: Mouse, Human 交差が予測される動物種: Rat, Guinea pig, Cow, Dog, Pig, Rhesus monkey, Gorilla, Elephant 
免疫原	Synthetic peptide, corresponding to a region within amino acids 1-50 of Human ZNF787 (NP_001002836.2).
ポジティブ・コントロール	HeLa, 293T, Jurkat and Mouse NIH3T3 whole cell lysates.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	pH: 6.8 Preservative: 0.09% Sodium azide Constituents: 99% Tris buffered saline, 0.1% BSA
精製度	Immunogen affinity purified
特記事項 (精製)	ab117837 was affinity purified using an epitope specific to ZNF787 immobilized on solid support.
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee Abpromise保証は、 次のテスト済みアプリケーションにおけるab117837の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
WB		1/2000 - 1/10000. Predicted molecular weight: 40 kDa.
IP		Use at 2-10 µg/mg of lysate.

ターゲット情報

機能	May be involved in transcriptional regulation.
配列類似性	Belongs to the krueppel C2H2-type zinc-finger protein family. Contains 7 C2H2-type zinc fingers.
翻訳後修飾	Phosphorylated upon DNA damage, probably by ATM or ATR.
細胞内局在	Nucleus.

画像



All lanes : Anti-ZNF787 antibody (ab117837) at 0.04 µg/ml

Lane 1 : HeLa whole cell lysate at 50 µg

Lane 2 : HeLa whole cell lysate at 15 µg

Lane 3 : 293T whole cell lysate at 50 µg


Lane 4 : Jurkat whole cell lysate at 50 µg

Lane 5 : NIH3T3 whole cell lysate at 50 µg

Developed using the ECL technique.

Predicted band size: 40 kDa

Exposure time: 3 minutes



Detection of Human ZNF787 by Immunoprecipitation in HeLa whole cell lysate (1 mg for IP, 20% of IP loaded) using ab117837 at 6 µg/mg of lysate. Subsequent WB detection was done using ab117837 at 1 µg/ml. ECL detection. Exposure time: 3 seconds.

Immunoprecipitation - Anti-ZNF787 antibody
(ab117837)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors