abcam

Product datasheet

Human PTHLH peptide ab42283

製品の詳細

製品名 Human PTHLH peptide

精製度 > 95 % HPLC.

Animal free No

由来 Synthetic

生物種 Human

配列 TRSAW

領域 107 to 111

特性

Our Abpromise guarantee covers the use of ab42283 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション Blocking

製品の状態 Lyophilized

前処理および保存

保存方法および安定性 Shipped at 4°C. After reconstitution store at -20°C. Avoid freeze / thaw cycles.

Constituent: 0.1% Acetic acid

再構成 Store at 2 to -8°C lyophilised, resuspend in 0.1% actetic acid

関連情報

機能 Neuroendocrine peptide which is a critical regulator of cellular and organ growth, development,

migration, differentiation and survival and of epithelial calcium ion transport. Regulates endochondral bone development and epithelial-mesenchymal interactions during the formation of the mammary glands and teeth. Required for skeletal homeostasis. Promotes mammary mesenchyme differentiation and bud outgrowth by modulating mesenchymal cell responsiveness to BMPs. Upregulates BMPR1A expression in the mammary mesenchyme and this increases the

sensitivity of these cells to BMPs and allows them to respond to BMP4 in a paracrine and/or autocrine fashion. BMP4 signaling in the mesenchyme, in turn, triggers epithelial outgrowth and

1

augments MSX2 expression, which causes the mammary mesenchyme to inhibit hair follicle formation within the nipple sheath (By similarity). Promotes colon cancer cell migration and invasion in an integrin alpha-6/beta-1-dependent manner through activation of Rac1.

Osteostatin is a potent inhibitor of osteoclastic bone resorption.

組織特異性 Ubiquitous. Also expressed in the mammary gland.

関連疾患 Brachydactyly E2

配列類似性 Belongs to the parathyroid hormone family.

翻訳後修飾 There are 3 principal secretory forms, called PTHrP[1-36], PTHrP[38-94], and osteostatin

(PTHrP[107-139]) arising from endoproteolytic cleavage of the initial translation product. Each of these secretory forms is believed to have one or more of its own receptors that mediates the

normal paracrine, autocrine and endocrine actions.

細胞内局在 Cytoplasm. Nucleus. Secreted.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors