abcam

Product datasheet

Human Versican peptide ab39784

製品の詳細

製品名 Human Versican peptide

精製度 > 90 % SDS-PAGE.

This peptide is greater than 70% pure.

Animal free No

由来 Synthetic

アミノ酸配列 1

生物種 Human

配列 CGG-DPEAAE

領域 436 to 441

アミノ酸配列2

生物種 Human

配列 CGG-DPEAAE

領域 436 to 441

特性

Our **Abpromise guarantee** covers the use of **ab39784** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション Blocking - Blocking peptide for Anti-Versican antibody (ab19345)

Neutralising

製品の状態 Lyophilized

前処理および保存

保存方法および安定性 Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

Double distilled water or equivalent after reconstitution.

再構成 Reconstitute with 0.1 ml of distilled water.

関連情報

1

機能 May play a role in intercellular signaling and in connecting cells with the extracellular matrix. May

take part in the regulation of cell motility, growth and differentiation. Binds hyaluronic acid.

組織特異性 Cerebral white matter and plasma. Isoform V0 and isoform V1 are expressed in normal brain,

gliomas, medulloblastomas, schwannomas, neurofibromas, and meningiomas. Isoform V2 is

restricted to normal brain and gliomas. Isoform V3 is found in all these tissues except

medulloblastomas.

関連疾患 Defects in VCAN are the cause of Wagner syndrome type 1 (WGN1) [MIM:143200]. WGN is a

dominantly inherited vitreoretinopathy characterized by an optically empty vitreous cavity with fibrillary condensations and a preretinal avascular membrane. Other optical features include progressive chorioretinal atrophy, perivascular sheating, subcapsular cataract and myopia.

Systemic manifestations are absent in WGN.

配列類似性 Belongs to the aggrecan/versican proteoglycan family.

Contains 1 C-type lectin domain. Contains 2 EGF-like domains.

Contains 1 lg-like V-type (immunoglobulin-like) domain.

Contains 2 Link domains.

Contains 1 Sushi (CCP/SCR) domain.

発生段階 Disappears after the cartilage development.

翻訳後修飾 Phosphorylation sites are present in the extracellular medium.

細胞内局在 Secreted > extracellular space > extracellular matrix.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors