abcam

Product datasheet

Mouse LYVE1 peptide ab34093

画像数 1

製品の詳細

製品名 Mouse LYVE1 peptide

精製度 > 90 % HPLC.

Animal free No

由来 Synthetic

生物種 Mouse

特性

Our Abpromise guarantee covers the use of ab34093 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

製品の状態

備考

Liquid

- First try to dissolve a small amount of peptide in either water or buffer. The more charged residues on a peptide, the more soluble it is in aqueous solutions.
- If the peptide doesn't dissolve try an organic solvent e.g. DMSO, then dilute using water or buffer.
- Consider that any solvent used must be compatible with your assay. If a peptide does not dissolve and you need to recover it, lyophilise to remove the solvent.
- Gentle warming and sonication can effectively aid peptide solubilisation. If the solution is cloudy or has gelled the peptide may be in suspension rather than solubilised.
- Peptides containing cysteine are easily oxidised, so should be prepared in solution just prior to use.

前処理および保存

保存方法および安定性

Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw

cycles.

Information available upon request.

関連情報

機能 Ligand-specific transporter trafficking between intracellular organelles (TGN) and the plasma

membrane. Plays a role in autocrine regulation of cell growth mediated by growth regulators containing cell surface retention sequence binding (CRS). May act as a hyaluronan (HA) transporter, either mediating its uptake for catabolism within lymphatic endothelial cells

 $themselves, or its \ transport\ into\ the\ lumen\ of\ afferent\ lymphatic\ vessels\ for\ subsequent\ re-uptake$

and degradation in lymph nodes.

組織特異性 Mainly expressed in endothelial cells lining lymphatic vessels.

配列類似性 Contains 1 Link domain.


翻訳後修飾 O-glycosylated.

細胞内局在 Membrane. Localized to the plasma membrane and in vesicles near extranuclear membranes

which may represent trans-Golgi network (TGN) and endosomes/prelysosomeal compartments.

Undergoes ligand-dependent internalization and recycling at the cell surface.

画像



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- · Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors