

Product datasheet

Mouse PSD95 peptide ab18661

2 References

製品の詳細

製品名	Mouse PSD95 peptide
精製度	> 90 % HPLC.
アクセッション番号	<u>Q62108-2</u>
Animal free	No
由来	Synthetic
生物種	Mouse

特性

Our **Abpromise guarantee** covers the use of **ab18661** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション Blocking - Blocking peptide for Anti-PSD95 antibody - Synaptic Marker (**ab18258**)

製品の状態 Liquid

備考

- First try to dissolve a small amount of peptide in either water or buffer. The more charged residues on a peptide, the more soluble it is in aqueous solutions.
- If the peptide doesn't dissolve try an organic solvent e.g. DMSO, then dilute using water or buffer.
- Consider that any solvent used must be compatible with your assay. If a peptide does not dissolve and you need to recover it, lyophilise to remove the solvent.
- Gentle warming and sonication can effectively aid peptide solubilisation. If the solution is cloudy or has gelled the peptide may be in suspension rather than solubilised.
- Peptides containing cysteine are easily oxidised, so should be prepared in solution just prior to use.

前処理および保存

保存方法および安定性 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.

Information available upon request.

関連情報

機能	Interacts with the cytoplasmic tail of NMDA receptor subunits and shaker-type potassium channels. Required for synaptic plasticity associated with NMDA receptor signaling. Overexpression or depletion of DLG4 changes the ratio of excitatory to inhibitory synapses in hippocampal neurons. May reduce the amplitude of ASIC3 acid-evoked currents by retaining the channel intracellularly. May regulate the intracellular trafficking of ADR1B.
組織特異性	Brain.
配列類似性	Belongs to the MAGUK family. Contains 1 guanylate kinase-like domain. Contains 3 PDZ (DHR) domains. Contains 1 SH3 domain.
ドメイン	The PDZ domain 3 mediates interaction with ADR1B. The L27 domain near the N-terminus of isoform 2 is required for HGS/HRS-dependent targeting to postsynaptic density.
翻訳後修飾	Palmitoylation of isoform 1 is required for targeting to postsynaptic density.
細胞内局在	Cell membrane. Cell junction, synapse, postsynaptic cell membrane, postsynaptic density. Cell projection, axon. Cell junction, synapse. High levels in postsynaptic density of neurons in the forebrain. Also in presynaptic region of inhibitory synapses formed by cerebellar basket cells on axon hillocks of Purkinje cells.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors