abcam

Product datasheet

Native Rabbit Fibrinogen protein ab92829

製品の詳細

製品名 Native Rabbit Fibrinogen protein

精製度 > 95 % SDS-PAGE.

Prepared from fresh rabbit plasma using several chromatographic steps. Plasminogen depleted

by lysine affinity chromatography.

発現系 Native

タンパク質長 Full length protein

Animal free No.

由来 Native

生物種 Rabbit

特性

Our **Abpromise guarantee** covers the use of **ab92829** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション SDS-PAGE

製品の状態 Liquid

備考 Thaw at 37°C without agitation until completely liquid, then gently mix before use. Keep fibrinogen

at 25-37°C, aliquot and flash freeze unused portion.

前処理および保存

保存方法および安定性 Shipped on dry ice. Upon delivery aliquot and store at -80°C. Avoid freeze / thaw cycles.

pH: 7.40

Constituent: 0.588% Sodium citrate

関連情報

機能 Fibrinogen has a double function: yielding monomers that polymerize into fibrin and acting as a

cofactor in platelet aggregation.

組織特異性 Plasma.

関連疾患 Defects in FGA are a cause of congenital afibrinogenemia (CAFBN) [MIM:202400]. This is a rare

1

autosomal recessive disorder characterized by bleeding that varies from mild to severe and by complete absence or extremely low levels of plasma and platelet fibrinogen. Note=The majority of cases of afibrinogenemia are due to truncating mutations. Variations in position Arg-35 (the site of cleavage of fibrinopeptide a by thrombin) leads to alpha-dysfibrinogenemias.

Defects in FGA are a cause of amyloidosis type 8 (AMYL8) [MIM:105200]; also known as systemic non-neuropathic amyloidosis or Ostertag-type amyloidosis. AMYL8 is a hereditary generalized amyloidosis due to deposition of apolipoprotein A1, fibrinogen and lysozyme amyloids. Viscera are particularly affected. There is no involvement of the nervous system. Clinical features include renal amyloidosis resulting in nephrotic syndrome, arterial hypertension, hepatosplenomegaly, cholestasis, petechial skin rash.

配列類似性 Contains 1 fibringen C-terminal domain.

A long coiled coil structure formed by 3 polypeptide chains connects the central nodule to the C-

terminal domains (distal nodules). The long C-terminal ends of the alpha chains fold back,

contributing a fourth strand to the coiled coil structure.

翻訳後修飾 The alpha chain is not glycosylated.

Forms F13A-mediated cross-links between a glutamine and the epsilon-amino group of a lysine

residue, forming fibronectin-fibrinogen heteropolymers.

About one-third of the alpha chains in the molecules in blood were found to be phosphorylated. Conversion of fibrinogen to fibrin is triggered by thrombin, which cleaves fibrinopeptides A and B from alpha and beta chains, and thus exposes the N-terminal polymerization sites responsible for the formation of the soft clot. The soft clot is converted into the hard clot by factor XIIIA which catalyzes the epsilon-(gamma-glutamyl)lysine cross-linking between gamma chains (stronger)

and between alpha chains (weaker) of different monomers.

Phosphorylation sites are present in the extracellular medium.

細胞内局在 Secreted.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors