

# Recombinant human BAFF-R protein ab50233

## 製品の詳細

製品名	Recombinant human BAFF-R protein
生理活性	Determined by its ability to block BAFF induced mouse splenocyte survival. The expected ED <sub>50</sub> for this effect is 2.0-4.0 µg/ml in the presence of 1.0 µg/ml of human soluble BAFF.
精製度	> 95 % SDS-PAGE.
発現系	Escherichia coli
タンパク質長	Protein fragment
Animal free	No
由来	Recombinant
生物種	Human
配列	MRRGPRSLRG RDAPAPTPCV PAECFDLLVR HCVACGLLRT PRPKPAGASS PAPRTALQPQ ESVGAGAGEA ALPLPG
領域	1 to 76

## 特性

Our **Abpromise guarantee** covers the use of **ab50233** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	SDS-PAGE Inhibition Assay
製品の状態	Lyophilized

## 前処理および保存

保存方法および安定性	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. pH: 4.00 Constituent: 0.129% Sodium citrate  Endotoxin level is less than 0.1 ng per µg (1EU/µg). This product is an active protein and may elicit a biological response in vivo, handle with caution.
------------	---

<b>再構成</b>	Centrifuge the vial prior to opening. Reconstitute in water to a concentration of 0.1-1.0 mg/ml. This solution can then be diluted into other aqueous buffers and stored at 4oC for 1 week or -20oC for future use
<b>関連情報</b>	
<b>機能</b>	B-cell receptor specific for TNFSF13B/TALL1/BAFF/BLyS. Promotes the survival of mature B-cells and the B-cell response.
<b>組織特異性</b>	Highly expressed in spleen and lymph node, and in resting B-cells. Detected at lower levels in activated B-cells, resting CD4+ T-cells, in thymus and peripheral blood leukocytes.
<b>関連疾患</b>	Defects in TNFRSF13C are the cause of immunodeficiency common variable type 4 (CVID4) [MIM:613494]; also called antibody deficiency due to BAFFR defect. CVID4 is a primary immunodeficiency characterized by antibody deficiency, hypogammaglobulinemia, recurrent bacterial infections and an inability to mount an antibody response to antigen. The defect results from a failure of B-cell differentiation and impaired secretion of immunoglobulins; the numbers of circulating B cells is usually in the normal range, but can be low.
<b>配列類似性</b>	Contains 1 TNFR-Cys repeat.
<b>細胞内局在</b>	Membrane.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors