abcam

Product datasheet

Recombinant human CDK6 + CCND3 protein ab84557

1 References 画像数 5

製品の詳細

製品名 Recombinant human CDK6 + CCND3 protein

生理活性 Specific activity of ab84557 was determined to be 36 nmol/min/mg.

精製度 > 75 % Densitometry.

Affinity purified.

発現系 Baculovirus infected Sf9 cells

タンパク質長 Full length protein

Animal free No

由来 Recombinant

生物種 Human

サブ His tag N-Terminus

特性

Our Abpromise guarantee covers the use of ab84557 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション Western blot

Functional Studies

SDS-PAGE

製品の状態 Liquid

備考 <u>ab56270</u> (Human Rb protein fragment) can be utilized as a substrate for assessing Kinase

activity

前処理および保存

保存方法および安定性 Shipped on dry ice. Upon delivery aliquot and store at -80°C. Avoid freeze / thaw cycles.

pH: 7.00

Preservative: 1.02% Imidazole

Constituents: 0.00174% PMSF, 0.82% Sodium phosphate, 0.00308% DTT, 25% Glycerol

(glycerin, glycerine), 1.74% Sodium chloride

This product is an active protein and may elicit a biological response in vivo, handle with caution.

1

関連情報


関連性

CDK family members are highly similar to the gene products of Saccharomyces cerevisiae cdc28, and Schizosaccharomyces pombe cdc2, and are known to be important regulators of cell cycle progression. This kinase is a catalytic subunit of the protein kinase complex that is important for cell cycle G1 phase progression and G1/S transition. The activity of Cdk6 first appears in mid-G1 phase, and is controlled by regulatory subunits including D-type cyclins and members of INK4 family of CDK inhibitors. This kinase, as well as CDK4, has been shown to phosphorylate, and thus regulate the activity of, tumor suppressor protein Rb. Cyclin D3 is a G1 cyclin closely related to cyclin D1 and D2. Cyclins D1, D2, and D3 are members of a growing superfamily of cyclins, proteins that govern transitions through distinct phases of the cell cycle by regulating the activity of cyclin-dependent kinases (cdks). All three known members of the D-type subfamily form active kinase complexes in vivo with a distinct subset of cdks, particularly with cdk4 and cdk6.


細胞内局在

CCND3: Cytoplasm. Nucleus

画像




The specific activity of CDK6 + CCND3 (ab84557) was determined to be 40 nmol/min/mg as per activity assay protocol




protein (ab84557)


SDS PAGE analysis of ab84557



SDS PAGE analysis of ab84557



SDS-PAGE of ab84557. CDK6 approximate MW 40kDa and CCND3 approximate MW 35kDa.



Kinase Assay: Specific activity of CDK6/CCND3 was determined to be 36 nmol/min/mg using protein substrate Rb.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors