

### Recombinant Human CSB protein ab114506

画像数 1

#### 製品の詳細

製品名	Recombinant Human CSB protein
発現系	Wheat germ
アクセッション番号	<b><u>Q03468-1</u></b>
タンパク質長	Protein fragment
Animal free	No
由来	Recombinant
生物種	Human
配列	RARNHLILPERLESESGHLQEASALLPTTEHDDLLVEMRNFI AFQAHTDG QASTREILQEFESKLSASQSCVFRELLRNLCFHRITSGGEGI WKLKPEYC
予測される分子量	37 kDa including tags
領域	1394 to 1493

#### 特性

Our **Abpromise guarantee** covers the use of **ab114506** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	ELISA
	SDS-PAGE
	Western blot
製品の状態	Liquid


#### 前処理および保存

保存方法および安定性	Shipped on dry ice. Upon delivery aliquot and store at -80°C. Avoid freeze / thaw cycles. pH: 8.00 Constituents: 0.3% Glutathione, 0.79% Tris HCl
------------	---

#### 関連情報

機能	Essential factor involved in transcription-coupled nucleotide excision repair which allows RNA polymerase II-blocking lesions to be rapidly removed from the transcribed strand of active genes. Upon DNA-binding, it locally modifies DNA conformation by wrapping the DNA around itself, thereby modifying the interface between stalled RNA polymerase II and DNA. It is required for transcription-coupled repair complex formation. It recruits the CSA complex (DCX(ERCC8) complex), nucleotide excision repair proteins and EP300 to the sites of RNA polymerase II-blocking lesions.
関連疾患	<p>Defects in ERCC6 are the cause of Cockayne syndrome type B (CSB) [MIM:133540]. Cockayne syndrome is a rare disorder characterized by cutaneous sensitivity to sunlight, abnormal and slow growth, cachectic dwarfism, progeroid appearance, progressive pigmentary retinopathy and sensorineural deafness. There is delayed neural development and severe progressive neurologic degeneration resulting in mental retardation. Two clinical forms are recognized: in the classical form or Cockayne syndrome type 1, the symptoms are progressive and typically become apparent within the first few years of life; the less common Cockayne syndrome type 2 is characterized by more severe symptoms that manifest prenatally. Cockayne syndrome shows some overlap with certain forms of xeroderma pigmentosum. Unlike xeroderma pigmentosum, patients with Cockayne syndrome do not manifest increased freckling and other pigmentation abnormalities in the skin and have no significant increase in skin cancer.</p> <p>Defects in ERCC6 are the cause of cerebro-oculo-facio-skeletal syndrome type 1 (COFS1) [MIM:214150]; also known as COFS syndrome or Pena-Shokeir syndrome type 2. COFS is a degenerative autosomal recessive disorder of prenatal onset affecting the brain, eye and spinal cord. After birth, it leads to brain atrophy, hypoplasia of the corpus callosum, hypotonia, cataracts, microcornea, optic atrophy, progressive joint contractures and growth failure. Facial dysmorphism is a constant feature. Abnormalities of the skull, eyes, limbs, heart and kidney also occur.</p> <p>Defects in ERCC6 are a cause of De Sanctis-Cacchione syndrome (DSC) [MIM:278800]; also known as xerodermic idiocy. DSC is an autosomal recessive syndrome consisting of xeroderma pigmentosum associated with mental retardation, retarded growth, gonadal hypoplasia and sometimes neurologic complications.</p> <p>Note=A genetic variation in the 5-prime flanking region of ERCC6 has been shown to be associated with susceptibility to age-related macular degeneration.</p> <p>Defects in ERCC6 are a cause of UV-sensitive syndrome (UVS) [MIM:600630]. UVS is a rare autosomal recessive disorder characterized by photosensitivity and mild freckling but without neurological abnormalities or skin tumors.</p>
配列類似性	<p>Belongs to the SNF2/RAD54 helicase family.</p> <p>Contains 1 helicase ATP-binding domain.</p> <p>Contains 1 helicase C-terminal domain.</p>
ドメイン	A C-terminal ubiquitin-binding domain (UBD) is essential for transcription-coupled nucleotide excision repair to proceed.
翻訳後修飾	<p>Phosphorylated upon DNA damage, probably by ATM or ATR.</p> <p>Ubiquitinated at the C-terminus. Ubiquitination by the CSA complex leads to ERCC6 proteasomal degradation in a UV-dependent manner.</p>
細胞内局在	Nucleus.

## 画像



SDS-PAGE - Recombinant Human CSB protein  
(ab114506)

ab114506 analysed by 12.5% SDS-PAGE and stained with Coomassie Blue.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

#### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors