

# Recombinant Human KAT13A / SRC1 protein ab82062

### 製品の詳細

---

|             |  |
|-------------|--|
| 製品名         | Recombinant Human KAT13A / SRC1 protein  |
| 精製度         | > 95 % SDS-PAGE.<br>ab82062 is purified by an affinity column.   |
| 発現系         | Escherichia coli   |
| アクセッション番号   | <b><u>Q15788</u></b>   |
| タンパク質長      | Protein fragment   |
| Animal free | No   |
| 由来          | Recombinant  |
| 生物種         | Human  |
| 領域          | 627 to 786   |
| タグ          | His tag N-Terminus   |
| 配列の追加情報     | This protein contains the Tobacco Etch virus (TEV) protease site & can be cleaved using acTEV to yield untagged SRC1. This protein will be useful for assays where a native untagged protein is desired. |

### 特性

---

Our **Abpromise guarantee** covers the use of **ab82062** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

|          |  |
|----------|--|
| アプリケーション | SDS-PAGE   |
| 製品の状態    | Liquid   |
| 備考       | ab82062 contains the Tobacco Etch virus (TEV) protease site and can be cleaved using acTEV to yield untagged KAT13A / SRC1 receptor interacting domain (RID). This protein will be useful for assays where a native untagged protein is desired.<br><br>ab82062 can also be used for protein-protein interaction assays. 100 units are sufficient for a protein-protein interaction assay. 1 unit equals 1 nanogram of purified protein. |

### 前処理および保存

---

|            |  |
|------------|--|
| 保存方法および安定性 | Shipped on dry ice. Upon delivery aliquot and store at -80°C. Avoid freeze / thaw cycles.<br><br>pH: 7.9<br>Constituents: 0.75% Potassium chloride, 0.0154% DTT, 0.316% Tris HCl, 0.00584% EDTA, 20% |
|------------|--|

## 関連情報

---

|       |  |
|-------|--|
| 機能    | Nuclear receptor coactivator that directly binds nuclear receptors and stimulates the transcriptional activities in a hormone-dependent fashion. Involved in the coactivation of different nuclear receptors, such as for steroids (PGR, GR and ER), retinoids (RXRs), thyroid hormone (TRs) and prostanoids (PPARs). Also involved in coactivation mediated by STAT3, STAT5A, STAT5B and STAT6 transcription factors. Displays histone acetyltransferase activity toward H3 and H4; the relevance of such activity remains however unclear. Plays a central role in creating multisubunit coactivator complexes that act via remodeling of chromatin, and possibly acts by participating in both chromatin remodeling and recruitment of general transcription factors. Required with NCOA2 to control energy balance between white and brown adipose tissues. Required for mediating steroid hormone response. Isoform 2 has a higher thyroid hormone-dependent transactivation activity than isoform 1 and isoform 3. |
| 組織特異性 | Widely expressed.  |
| 関連疾患  | Note=A chromosomal aberration involving NCOA1 is a cause of rhabdomyosarcoma. Translocation t(2;2)(q35;p23) with PAX3 generates the NCOA1-PAX3 oncogene consisting of the N-terminus part of PAX3 and the C-terminus part of NCOA1. The fusion protein acts as a transcriptional activator. Rhabdomyosarcoma is the most common soft tissue carcinoma in childhood, representing 5-8% of all malignancies in children.   |
| 配列類似性 | Belongs to the SRC/p160 nuclear receptor coactivator family.<br>Contains 1 basic helix-loop-helix (bHLH) domain.<br>Contains 1 PAS (PER-ARNT-SIM) domain.  |
| ドメイン  | The C-terminal (1107-1441) part mediates the histone acetyltransferase (HAT) activity. Contains 7 Leu-Xaa-Xaa-Leu-Leu (LXXLL) motifs. LXXLL motifs 3, 4 and 5 are essential for the association with nuclear receptors. LXXLL motif 7, which is not present in isoform 2, increases the affinity for steroid receptors in vitro.   |
| 翻訳後修飾 | Sumoylated; sumoylation increases its interaction with PGR and prolongs its retention in the nucleus. It does not prevent its ubiquitination and does not exert a clear effect on the stability of the protein.<br>Ubiquitinated; leading to proteasome-mediated degradation. Ubiquitination and sumoylation take place at different sites.<br>Phosphorylated upon DNA damage, probably by ATM or ATR.   |
| 細胞内局在 | Nucleus.   |

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors