

Recombinant Human MDM2 protein ab82080

3 References [画像数 1](#)

製品の詳細

製品名	Recombinant Human MDM2 protein
精製度	> 95 % SDS-PAGE. ab82080 is purified by affinity and FPLC chromatography and is greater than 95% homogeneous based on SDS-PAGE analysis.
発現系	Escherichia coli
アクセッション番号	<u>Q00987</u>
タンパク質長	Full length protein
Animal free	No
由来	Recombinant
生物種	Human
配列	MAHHHHHHASMCNTNMSVPTDGAVTTSQIPASEQETLVRPKP LLLKLLKS VGAQKDTYTMKEVLFYLGQYIMTKRLYDEKQQHIVYCSNDLL GDLFGVPS FSVKEHRKIYTMIRNLVVVNQQESSDSGTSVSENCHLEGG SDQKDLVQ ELQEEKPSSSHLVSRPSTSSRRRAISETEENSDELSEGERQRK RHKSDSIS LSFDESLALCVIREICCERSSSSESTGTPSNPDL DAGVSEHS GDWLDQDS VSDQFSVEFEVESL DSEYSLSEEGQEL SDEDDEVYQVTVYQ AGESDTDS FEEDPEISLADYWKCTSCNEMNPPLPSHCNRCWALRENWLPE DKGKDKGE ISEKAKLENSTQAEEGFDVPDCKKTIVNDSRESCVEENDDKI TQASQSQE SEDYSQPSTSSSIIYSSQEDVKEFEREETQDKEESVESSLPL NAIEPCVI CQGRPKNGCIVHGKTGHLMACFTCAKLLKRNKPCPVCRQPI QMIVLTYF PGLHHHHHHHH
予測される分子量	55 kDa
実際の分子量	58 kDa

特性

Our **Abpromise guarantee** covers the use of **ab82080** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	SDS-PAGE
製品の状態	Liquid
備考	1 unit equals 1 nanogram of purified protein.

前処理および保存


保存方法および安定性	Shipped on dry ice. Upon delivery aliquot and store at -80°C. Avoid freeze / thaw cycles. pH: 7.9 Constituents: 0.75% Potassium chloride, 0.0154% DTT, 0.316% Tris HCl, 0.00584% EDTA, 20% Glycerol
------------	---

関連情報

機能	E3 ubiquitin-protein ligase that mediates ubiquitination of p53/TP53, leading to its degradation by the proteasome. Inhibits p53/TP53- and p73/TP73-mediated cell cycle arrest and apoptosis by binding its transcriptional activation domain. Also acts as an ubiquitin ligase E3 toward itself and ARRB1. Permits the nuclear export of p53/TP53. Promotes proteasome-dependent ubiquitin-independent degradation of retinoblastoma RB1 protein. Inhibits DAXX-mediated apoptosis by inducing its ubiquitination and degradation. Component of the TRIM28/KAP1-MDM2-p53/TP53 complex involved in stabilizing p53/TP53. Also component of the TRIM28/KAP1-ERBB4-MDM2 complex which links growth factor and DNA damage response pathways.
組織特異性	Ubiquitous. Isoform Mdm2-A, isoform Mdm2-B, isoform Mdm2-C, isoform Mdm2-D, isoform Mdm2-E, isoform Mdm2-F and isoform Mdm2-G are observed in a range of cancers but absent in normal tissues.
関連疾患	Note=Seems to be amplified in certain tumors (including soft tissue sarcomas, osteosarcomas and gliomas). A higher frequency of splice variants lacking p53 binding domain sequences was found in late-stage and high-grade ovarian and bladder carcinomas. Four of the splice variants show loss of p53 binding.
配列類似性	Belongs to the MDM2/MDM4 family. Contains 1 RanBP2-type zinc finger. Contains 1 RING-type zinc finger. Contains 1 SWIB domain.
ドメイン	Region I is sufficient for binding p53 and inhibiting its G1 arrest and apoptosis functions. It also binds p73 and E2F1. Region II contains most of a central acidic region required for interaction with ribosomal protein L5 and a putative C4-type zinc finger. The RING finger domain which coordinates two molecules of zinc interacts specifically with RNA whether or not zinc is present and mediates the heterooligomerization with MDM4. It is also essential for its ubiquitin ligase E3 activity toward p53 and itself.
翻訳後修飾	Phosphorylated in response to ionizing radiation in an ATM-dependent manner. Auto-ubiquitinated; which leads to proteasomal degradation. Deubiquitinated by USP2 leads to its accumulation and increases deubiquitination and degradation of p53/TP53. Deubiquitinated by USP7; leading to stabilize it.
細胞内局在	Nucleus > nucleoplasm. Cytoplasm. Nucleus > nucleolus. Expressed predominantly in the

nucleoplasm. Interaction with ARF(P14) results in the localization of both proteins to the nucleolus. The nucleolar localization signals in both ARF(P14) and MDM2 may be necessary to allow efficient nucleolar localization of both proteins. Colocalizes with RASSF1 isoform A in the nucleus.

画像



SDS-PAGE analysis of Recombinant Human MDM2 protein (ab82080).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors