

# Recombinant human NEK2 protein ab42599

## 製品の詳細

製品名	Recombinant human NEK2 protein
生理活性	100 U/mg. One unit is defined as the amount of enzyme that will phosphorylate 1 nmol of myelin basic protein (MBP) substrate per minute at pH 7.4 and 30°C. Assay buffer: 50 mM HEPES, pH 7.4, 3 mM MgCl <sub>2</sub> , 3 mM MnCl <sub>2</sub> , 1 mM DTT, 3 uM Naorthovanadate, 0.5 mM ATP, 0.3 mg/ml MBP substrate, and 0.2 ug/ml NEK2.
精製度	> 70 % SDS-PAGE. Affinity purified.
発現系	Baculovirus infected Sf9 cells
タンパク質長	Full length protein
Animal free	No
由来	Recombinant
生物種	Human

## 特性

Our **Abpromise guarantee** covers the use of **ab42599** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Inhibition Assay
製品の状態	Liquid
備考	Expressed in a Baculovirus infected Sf9 cell expression system.

## 前処理および保存

保存方法および安定性	Shipped on Dry Ice. Upon delivery aliquot. Store at -80°C. Avoid freeze / thaw cycle. pH: 8.00 Constituents: 0.0462% (R*,R*)-1,4-Dimercaptobutan-2,3-diol, 0.395% Tris HCl, 0.05% Tween, 40% Glycerol (glycerin, glycerine), 0.87% Sodium chloride This product is an active protein and may elicit a biological response in vivo, handle with caution.
------------	--

## 関連情報

<b>機能</b>	Protein kinase that is involved in mitotic regulation. Integral component of the mitotic spindle-assembly checkpoint which is necessary for proper chromosome segregation during metaphase-anaphase transition. Required for association of MAD2L1 to kinetochore. Phosphorylates SGOL1. May have a role at the G2-M transition. May also play a role in meiosis. Isoform 1 but not isoform 2 appears to play a role in centrosome splitting. Isoform 1 phosphorylates and activates NEK11 in G1/S-arrested cells. Isoform 2, which is not present in the nucleolus, does not.
<b>組織特異性</b>	Isoform 1 and isoform 2 are expressed in peripheral blood T-cells and a wide variety of transformed cell types.
<b>配列類似性</b>	Belongs to the protein kinase superfamily. NEK Ser/Thr protein kinase family. NIMA subfamily. Contains 1 protein kinase domain.
<b>発生段階</b>	Accumulates throughout S phase and shows maximal levels in late G2. This expression pattern is highly reminiscent of that of A and B cyclins. Isoform 1 and isoform 2 expression is low in the G1 phase and increases in the S/G2 phases. Isoform 1 is absent from cells arrested in the G2/M prometaphase. Isoform 2 remains present in cells arrested in the G2/M prometaphase.
<b>翻訳後修飾</b>	It is unsure whether Thr-170 or Ser-171 is phosphorylated.
<b>細胞内局在</b>	Cytoplasm. Predominantly cytoplasmic; Nucleus > nucleolus. Has a nucleolar targeting/ retention activity via a coiled-coil domain at the C-terminal end and Nucleus. Chromosome > centromere. Chromosome > centromere > kinetochore. Co-localizes with SGOL1 and MAD1L1 at the kinetochore. Not associated with kinetochore in the interphase but becomes associated with it upon the breakdown of the nuclear envelope.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors