abcam

Product datasheet

Recombinant human PKC epsilon protein ab60847

画像数5

製品の詳細

製品名 Recombinant human PKC epsilon protein

精製度 > 90 % Densitometry.

Affinity purified.

発現系 Baculovirus infected Sf9 cells

タンパク質長 Full length protein

Animal free No

由来 Recombinant

生物種 Human

タブ GST tag N-Terminus

特性

Our <u>Abpromise guarantee</u> covers the use of ab60847 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション Functional Studies

SDS-PAGE Western blot

製品の状態 Liquid

備考 <u>ab204875</u> (PKC epsilon peptide) can be utilized as a substrate for assessing kinase activity

前処理および保存

保存方法および安定性 Shipped on dry ice. Upon delivery aliquot and store at -80°C. Avoid freeze / thaw cycles.

pH: 7.50

Constituents: 0.0038% EGTA, 1.74% PMSF, 0.00385% DTT, 0.79% Tris HCI, 0.00292% EDTA,

25% Glycerol (glycerin, glycerine), 0.87% Sodium chloride

This product is an active protein and may elicit a biological response in vivo, handle with caution.

関連情報

1

機能 This is calcium-independent, phospholipid-dependent, serine- and threonine-specific enzyme.

PKC is activated by diacylglycerol which in turn phosphorylates a range of cellular proteins. PKC

also serves as the receptor for phorbol esters, a class of tumor promoters.

配列類似性 Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. PKC subfamily.

Contains 1 AGC-kinase C-terminal domain.

Contains 1 C2 domain.

Contains 2 phorbol-ester/DAG-type zinc fingers.


Contains 1 protein kinase domain.

ドメイン The C1 domain, containing the phorbol ester/DAG-type region 1 (C1A) and 2 (C1B), is the

diacylglycerol sensor and the C2 domain is a non-calcium binding domain.


翻訳後修飾 Phosphorylation on Thr-566 by PDPK1 triggers autophosphorylation on Ser-729.

画像




Functional Studies - Recombinant human PKC epsilon protein (ab60847)

The specific activity of PKC epsilon (ab60847) was determined to be 1122 nmol/min/mg as per activity assay protocol and was equivalent to 1398 nmol/min/mg as per radiometric assay




protein (ab60847)


SDS PAGE analysis of ab60847



SDS PAGE analysis of ab60847



Sample Kinase Activity Plot.



ab60847 on SDS-PAGE, MW ~110 kDa.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors