

Recombinant Human Retinoid X Receptor alpha/RXRA protein ab84128

製品の詳細

製品名	Recombinant Human Retinoid X Receptor alpha/RXRA protein
生理活性	1 unit equals 1 nanogram purified protein.
精製度	> 95 % SDS-PAGE.
発現系	Escherichia coli
タンパク質長	Protein fragment
Animal free	No
由来	Recombinant
生物種	Human
領域	200 to 462

特性

Our **Abpromise guarantee** covers the use of **ab84128** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	SDS-PAGE
製品の状態	Liquid
備考	1 unit equals 1 nanogram purified protein. This product was previously labelled as Retinoid X Receptor alpha

前処理および保存

保存方法および安定性	Shipped on dry ice. Upon delivery aliquot and store at -80°C. Avoid freeze / thaw cycles. pH: 7.9 Constituents: 0.75% Potassium chloride, 0.0154% DTT, 0.316% Tris HCl, 0.00584% EDTA, 20% Glycerol (glycerin, glycerine)
------------	---

関連情報

機能	Receptor for retinoic acid. Retinoic acid receptors bind as heterodimers to their target response elements in response to their ligands, all-trans or 9-cis retinoic acid, and regulate gene expression in various biological processes. The RAR/RXR heterodimers bind to the retinoic acid response elements (RARE) composed of tandem 5'-AGGTCA-3' sites known as DR1-DR5. The high affinity ligand for RXRs is 9-cis retinoic acid. RXRA serves as a common heterodimeric partner for a number of nuclear receptors. The RXR/RAR heterodimers bind to the retinoic acid response elements (RARE) composed of tandem 5'-AGGTCA-3' sites known as DR1-DR5. In the absence of ligand, the RXR-RAR heterodimers associate with a multiprotein complex containing transcription corepressors that induce histone acetylation, chromatin condensation and transcriptional suppression. On ligand binding, the corepressors dissociate from the receptors and associate with the coactivators leading to transcriptional activation. The RXRA/PPARA heterodimer is required for PPARA transcriptional activity on fatty acid oxidation genes such as ACOX1 and the P450 system genes.
組織特異性	Highly expressed in liver, also found in lung, kidney and heart.
配列類似性	Belongs to the nuclear hormone receptor family. NR2 subfamily. Contains 1 nuclear receptor DNA-binding domain.
ドメイン	Composed of three domains: a modulating N-terminal domain (AF1 domain), a DNA-binding domain and a C-terminal ligand-binding domain (AF2 domain).
翻訳後修飾	Phosphorylated on serine and threonine residues mainly in the N-terminal modulating domain. Constitutively phosphorylated on Ser-21 in the presence or absence of ligand. Under stress conditions, hyperphosphorylated by activated JNK on Ser-56, Ser-70, Thr-82 and Ser-260 (By similarity). Phosphorylated on Ser-27, in vitro, by PKA. This phosphorylation is required for repression of cAMP-mediated transcriptional activity of RARA. Sumoylation negatively regulates transcriptional activity. Desumoylated specifically by SENP6.
細胞内局在	Nucleus.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors