

Recombinant human Serum Amyloid A (mutated D60N + H71R) protein ab50232

画像数 1

製品の詳細

製品名	Recombinant human Serum Amyloid A (mutated D60N + H71R) protein
生理活性	Tested by its ability to down- regulate lipid biosynthesis in aortic smooth muscle cells. The effective concentration was found to be 4 μ M. Schreiber, BM. et al. Biochem J. 1999 Nov 15; 344 Pt 1:7-13
精製度	>= 98 % SDS-PAGE. >=98% by SDS-PAGE and HPLC analyses.
エンドトキシン・レベル	< 0.100 Eu/ μ g
発現系	Escherichia coli
アクセッション番号	<u>P0DJ18</u>
タンパク質長	Full length protein
Animal free	No
由来	Recombinant
生物種	Human
配列	MRSFFSFLGE AFDGARDMWR AYSDMREANY IGSDKYFHAR GNYDAAKRGF GGVWAAEAIS NARENIQRFF GRGAEDSLAD QAANEWGRSG KDPNHFRPAG LPEKY
予測される分子量	12 kDa
修飾	mutated D60N + H71R

特性

Our **Abpromise guarantee** covers the use of **ab50232** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	SDS-PAGE Functional Studies Western blot
製品の状態	Lyophilized
備考	For most in vitro applications, apo-SAA exerts its biological activity in the concentration range of

10-50 µg/ml. ab50232 can be used as a WB positive control in conjunction with [ab687](#).

前処理および保存

保存方法および安定性

Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

This product is an active protein and may elicit a biological response in vivo, handle with caution.

再構成


Please reconstitute this product in water to a concentration of 1mg/ml.

関連情報

関連性

Function: Major acute phase reactant. Apolipoprotein of the HDL complex. Tissue specificity: Expressed by the liver; secreted in plasma. Disease: Note=Reactive, secondary amyloidosis is characterized by the extracellular accumulation in various tissues of the SAA protein. These deposits are highly insoluble and resistant to proteolysis; they disrupt tissue structure and compromise function. Note=Elevated serum SAA protein levels may be associated with lung cancer. Similarity: Belongs to the SAA family. PTM: This protein is the precursor of amyloid protein A, which is formed by the removal of approximately 24 residues from the C-terminal end.

画像



Anti-Serum Amyloid A antibody [115] ([ab687](#)) at 0.5 µg/ml +
Recombinant human Serum Amyloid A (mutated D60N + H71R)
protein (ab50232) at 0.01 µg

Secondary

Goat Anti-Mouse IgG H&L (HRP) preadsorbed ([ab97040](#)) at
1/5000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Exposure time: 1 minute

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors