

Recombinant human TIMP2 protein ab82117

1 References

製品の詳細

製品名	Recombinant human TIMP2 protein
生理活性	TIMP2 activity was measured by its ability to inhibit human MMP1 induced hydrolysis of a chromogenic peptide substrate at room temperature. Half maximal inhibition was obtained at a TIMP2 concentration of approximately 0.5 µg/ml, when using an MMP1 concentration of 1.6 µg/ml.
精製度	> 95 % SDS-PAGE. Purity is greater than 95% by SDS-PAGE gel and HPLC analyses. Endotoxin level is less than 0.1 ng per µg (1EU/µg).
発現系	Escherichia coli
アクセッション番号	P16035
タンパク質長	Full length protein
Animal free	No
由来	Recombinant
生物種	Human
配列	CSCSPVHPQ QAFCNADVVI RAKAVSEKEV DSGNDIYGNP IKRIQYEIKQ IKMFKGPEKD IEFIYTAPSS AVCGVSLDVG GKKEYLIAGK AEGDGKMHIT LCDFIVPWDT LSTTQKKSLN HRYQMGCECK ITRCPMIPCY ISSPDECLWM DWVTEKNING HQAFFACIK RSDGSCAWYR GAAPPKQEFL DIEDP
領域	2 to 220

特性

Our **Abpromise guarantee** covers the use of **ab82117** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	SDS-PAGE Functional Studies
製品の状態	Lyophilized
備考	Centrifuge the vial prior to opening. Reconstitute in water to a concentration of 0.1-1.0 mg/ml. Do not vortex (can be stored at 4C for up to 1 week). For extended storage, it is recommended to further dilute in a buffer containing a carrier protein

(example 0.1% BSA) and store in working aliquots at -20°C to -80°C.

前処理および保存

保存方法および安定性

Shipped at 4°C. The lyophilized protein is stable for a few weeks at room temperature. Store at -20°C or -80°C. Avoid freeze / thaw cycle. Please see notes section.

This product is an active protein and may elicit a biological response in vivo, handle with caution.

再構成

Centrifuge the vial prior to opening. Reconstitute in water to a concentration of 0.1-1.0 mg/ml. Do not vortex (can be stored at 4°C for up to 1 week). For extended storage, it is recommended to further dilute in a buffer containing a carrier protein (example 0.1% BSA) and store in working aliquots at -20°C to -80°C.

関連情報

機能

Complexes with metalloproteinases (such as collagenases) and irreversibly inactivates them by binding to their catalytic zinc cofactor. Known to act on MMP-1, MMP-2, MMP-3, MMP-7, MMP-8, MMP-9, MMP-10, MMP-13, MMP-14, MMP-15, MMP-16 and MMP-19.

配列類似性

Belongs to the protease inhibitor I35 (TIMP) family.
Contains 1 NTR domain.

翻訳後修飾

The activity of TIMP2 is dependent on the presence of disulfide bonds.

細胞内局在

Secreted.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors