abcam

Product datasheet

Recombinant Streptolysin O (Hemolytic streptococcus) protein ab63978

5 References 画像数 1

製品の詳細

製品名 Recombinant Streptolysin O (Hemolytic streptococcus) protein

生理活性 Definition of 1HU is activation of 50% hemolysis by incubating 3% sheep red blood cells at 37?

for 30 minutes.

The specific activity is as high as 1,200,000-2,000,000 hemolytic units (HU) /mg.

精製度 > 98 % SDS-PAGE.

Affinity-purified with Ni colum followed by gel-filtration

発現系 Escherichia coli

アクセッション番号 <u>Q54114</u>

タンパク質長 Full length protein

Animal free No.

由来 Recombinant

生物種 Streptococcus

配列 MGGSHHHHHH GMASMTGGQQ MGRDLYDDDD KDPSSR

NKQN TANTETTTTN EQPKPESSEL TTEKAGQKMD

DMLNSNDMIK LAPKEMPLES AEKEEKKSED
NKKSEEDHTE EINDKIYSLN YNELEVLAKN
GETIENFVPK EGVKKADKFI VIERKKKNIN
TTPVDISIID SVTDRTYPAA LQLANKGFTE

NDPTYANVST AIDNLVNQWH DNYSGGNTLP ARTQYTESMV YSKSQIEAAL NVNSKILDGT LGIDFKSISK GEKKVMIAAY KQIFYTVSAN

NKPDAVVTKR NPQKIHIDLP GMGDKATVEV

FVSNVAYGRT VFVKLETSSK SNDVEAAFSA ALKGTDVKTN GKYSDILENS SFTAVVLGGD

LPNNPADVFD KSVTLKELQR KGVSNEAPPL

AAEHNKVVTK DFDVIRNVIK DNATFSRKNP AYPISYTSVF LKNNKIAGVN NRSEYVETTS

TEYTSGKINL SHQGAYVAQY EILWDEINYD DKGKEVITKR RWDNNWYSKT SPFSTVIPLG

ANSRNIRIMA RECTGLAWEW WRKVIDERDV

1

領域 37 to 574

サブ His tag N-Terminus

特性

Our Abpromise guarantee covers the use of ab63978 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション SDS-PAGE

Functional Studies

製品の状態 Liquid

備考 Information about health hazards:

Target: Cholesterol on human and animal cell membrane

Health Hazards: May be fatal if enters bloodstream.

LD50-Lethal dose (50 percent kill) intravenous,

Rabbit, 1500 ng/kg (Ref :PHTHDT Pharmacology and Therapeutics. (Pergamon Press Ltd.,Headington Hill Hall, Oxford OX3 0BW, UK) Vol.(Issue) 11, Page 661,1981)

Guinea pig, 12 ug/kg (Ref: BICMBE Biochimie. (SPPIF, B.P.22, F-41353 Vineuil, France, Vol.

(Issue)55,Page 1187, 1973)

Toxicity is much less when introduced via other routes of entry like interdermal injection.

前処理および保存

保存方法および安定性

Shipped on Dry Ice. Store at -80°C. Avoid freeze / thaw cycle.

pH: 6

Constituents: 49% Water, 0.015% DTT, 50% Glycerol (glycerin, glycerine), 0.1% Streptolysin, x1

Phosphate Buffered Saline


This product is an active protein and may elicit a biological response in vivo, handle with caution.

関連情報

関連性

Streptolysin O (SLO) is a membrane-damaging extracellular toxin produced by hemolytic streptococci. The membrane-damaging activity is measured by hemolysis of red-blood cell. SLO is oxygen-sensitive and is easily inactivated in its presence but can be reactivated by thiol compounds , so it is also called thiol-activated cytolysin. SLO is produced not only by Group A hemolytic streptococci but also by Group C and Group G strains. The amino acid sequences are highly conserved among them and their homology is over 98%.

画像



SDS-PAGE analysis of purified Streptolysin O (Hemolytic streptococcus) protein (ab63978).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors