

Biotinylated Goat anti-Mouse & Rabbit IgG (H+L) (Ready To Use) ab64257

[10 References](#) [画像数 1](#)

製品の概要

製品名	Biotinylated Goat anti-Mouse & Rabbit IgG (H+L) (Ready To Use)
アプリケーション	適用あり: IHC-P
特記事項	<p>Staining protocol</p> <p>Deparaffinize and rehydrate formalin-fixed paraffin-embedded tissue section. Add enough drops of an appropriate hydrogen peroxide block to cover the sections. Incubate for 10 minutes. Wash 2 times in buffer. Perform appropriate pretreatment if required. Wash 3 times in buffer. Apply an appropriate protein block and incubate for 5 minutes at room temperature to block nonspecific background staining. Wash 1 time in buffer. Apply primary antibody and incubate according to manufacturer's protocol. Wash 4 times in buffer. Apply Biotinylated Goat anti Mouse & Rabbit IgG(H+L) solution (ab64257) and incubate for 10 minutes at room temperature. Wash 4 times in buffer. Apply enzyme labeled streptavidin and incubate according to manufacturer's protocol. Rinse 4 times in buffer. Place slide in appropriate chromogenic substrate and incubate until desired reaction is achieved. Rinse 4 times in buffer. Add enough drops of Hematoxylin to cover the section. Incubate for 1 minute. Rinse 7-8 times in tap water. Add Mounting Medium to cover the section.</p> <p>This reagent is intended to be used in a labeled streptavidin-biotin immunoenzymatic antigen detection system. This technique involves the sequential incubation of the specimen with an unconjugated primary antibody specific to the target antigen, a biotinylated secondary antibody which reacts with the primary antibody, enzyme-labeled streptavidin, and substrate-chromogen.</p>

製品の特性

製品の状態	Liquid
保存方法	Store at +4°C.
バッファー	pH: 7.2 Preservative: 0.1% Sodium azide Constituent: PBS
IHC 試薬 備考	This reagent is intended to be used in a labeled streptavidin-biotin immunoenzymatic antigen detection system. This technique involves the sequential incubation of the specimen with an unconjugated primary antibody specific to the target antigen, a biotinylated secondary antibody

which reacts with the primary antibody, enzyme-labeled streptavidin, and substrate-chromogen.

細胞内局在

Secreted

細胞内局在


Secreted

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab64257の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		Use at an assay dependent dilution.

画像



Human colon carcinoma stained with anti-p53 antibody using Biotinylated Goat anti Mouse & Rabbit IgG (H+L) (Ready To Use) (ab64257).

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Biotinylated Goat anti-Mouse & Rabbit IgG (H+L) (Ready To Use) (ab64257)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors