

Streptavidin HRP (ready-to-use) ab64269

★★★★★ [1 Abreviews](#) [42 References](#) [画像数 2](#)

製品の概要

製品名	Streptavidin HRP (ready-to-use)
アプリケーション	適用あり: IHC-P
特記事項	<p>Streptavidin-HRP for use with biotinylated secondary antibodies during IHC / immunohistochemistry. Optimized, ready-to-use format.</p> <p>This product was previously called Streptavidin-Peroxidase.</p> <p>IHC protocol suitable for use with Streptavidin-HRP ab64269:</p> <ol style="list-style-type: none"> 1. Deparaffinize and rehydrate formalin-fixed paraffin-embedded tissue section. 2. Use appropriate <u>antigen retrieval buffer or enzyme</u> (primary antibody dependent) to treat sections. Wash 3 times in buffer. 3. Add enough <u>hydrogen peroxide blocking solution</u> to cover the sections. Incubate for 10 minutes. Wash 2 times in buffer. If necessary, use <u>avidin biotin blocking</u>. 4. Apply <u>protein block</u> (or <u>normal serum</u> from same species as secondary antibody) and incubate for 5 minutes at room temperature to block nonspecific background staining. Wash once in buffer. 5. Apply primary antibody in <u>antibody diluent</u> and incubate. 6. Wash 4 times in buffer. Incubate slide with <u>biotinylated secondary antibody</u> (or <u>HRP polymer secondary antibody</u> and skip step 7). Wash 4 times in buffer. 7. Apply streptavidin-HRP ab64269 and incubate for 10 minutes at room temperature. 8. Rinse 4 times in buffer. Place slide in <u>DAB substrate</u> or <u>AEC substrate</u> and incubate until desired color is achieved. Rinse 4 times in buffer. 9. Add enough drops of <u>hematoxylin</u> to cover the section. Incubate for 1 minute. 10. Rinse 7-8 times in tap water. Add <u>mounting medium</u> to cover the section. <p>Find complete IHC kits, and reagents for antigen retrieval, blocking, signal amplification, visualization, counterstaining, and mounting in the <u>IHC kits and reagents guide</u>.</p>

製品の特性

製品の状態	Liquid
-------	--------


保存方法	Store at +4°C.
バッファー	pH: 7.60 Preservative: 0.2% Proclin 300 Constituent: PBS
関連性	Streptavidin peroxidase is used in a labeled streptavidin-biotin immunoenzymatic antigen detection system. This technique involves the sequential incubation of the specimen with an unconjugated primary antibody specific to the target antigen, a biotinylated secondary antibody which reacts with the primary antibody, enzyme-labeled streptavidin, and substrate-chromogen.

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab64269の使用に適用されます**
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
IHC-P		Use at an assay dependent concentration.

画像



ab64269 Straptavidin Peroxidase used along with an anti-CD20 antibody on Human Tonsil

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Streptavidin Peroxidase (Ready to Use) (ab64269)



Human colon carcinoma stained with anti-p53 antibody using Streptavidin Peroxidase (ab64269).

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Streptavidin Peroxidase (Ready to Use) (ab64269)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors