

Heparin Sepharose® ab193268

製品の概要

製品名	Heparin Sepharose®
アッセイタイプ	Bead-based sandwich immunoassay (quantitative, multiplexable)
検出感度	>= 0.4 mg/ml
製品の概要	High binding capacity (>0.4mg/mL). Minimal leaching of ligand. For column or batch purification of heparin-binding proteins & cation exchange (ab193268).

Contents:

Supplied as a 50% slurry in 20 % Ethanol; >2.5 mg heparin per mL Sepharose® beads.

Features:

Heparin beads have been widely used in affinity purification of various heparin-binding proteins or ligands, such as antithrombin III, lipoprotein, as well as DNA binding proteins (transcription factors, virus coat proteins etc). Abcam's Heparin Sepharose® is designed for purification of heparin-binding proteins and ligands. It can also be used as a high capacity cation exchange medium. Specific proteins can be separated by using different concentrations of salt or a salt gradient. This Heparin Sepharose® formulation exhibits excellent binding capacity, high flow rate, no significant loss of the heparin ligand and a pH stability range of 2-10.

These beads are for use in column purification. If used in batch purification, we recommend not exceeding 150 x g when centrifuging.

Store beads at 4°C.

The beads may be damaged above 40°C.

DO NOT FREEZE.

Wash beads 3 times with 3x bead volume of desired buffer before use.

Applications:

Purification of heparin-binding proteins, enzymes or other ligands.

Sepharose is a registered trademark of GE Healthcare

特記事項

This product is manufactured by BioVision, an Abcam company and was previously called 6553 Heparin Sepharose. 6553-10 is the same size as the 10 ml size of ab193268.

Heparin Sepharose[®] is prepared by covalently coupling heparin to epoxy-activated 6% cross-linked Sepharose[®] beads. The coupling was optimized to give a high binding capacity and could be greater than 0.4 mg of heparin-binding protein (such as thrombin) per ml of wet gel.

Suggested Protocol:

Wash column with ddH₂O to remove air bubbles.

Fill column with heparin beads.

Wash the column with 5X volume of Binding Buffer.

Dilute sample with Binding Buffer (1:1 ratio) or change the sample solution to binding buffer by means of your choice.

Add the sample solution onto the column.

Collect the solution and repeat step 5 & 6 several times if necessary.

Wash the column 5-10 times with the Binding Buffer.

Add Elution Buffer to elute bound protein.

Collect the eluent using microcentrifuge tube.

Assay protein concentration and combine the fractions containing sufficient heparin-binding protein.

Bead can be cleaned and regenerated by washing with 2-3x volume of high concentration salt solution and then the binding buffer.

アプリケーション

適用あり: Purification

製品の特徴

保存方法

Store at +4°C. Please refer to protocols.

内容	10 ml	1 ml	50 ml
Heparin Sepharose [®]	1 x 10ml	1 x 1ml	1 x 50ml

アプリケーション

The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab193268の使用に適用されます

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Purification		Use at an assay dependent concentration. Purification of heparin-binding proteins, enzymes or other ligands.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors