

Donkey Anti-Goat IgG H&L (DyLight® 488) preadsorbed ab96935

★★★★★ [1 Abreviews](#) [7 References](#) [画像数 3](#)

製品の概要

製品名	Donkey Anti-Goat IgG H&L (DyLight® 488) preadsorbed
由来種	Donkey
ターゲット生物種	Goat
特異性	By immunoelectrophoresis and ELISA this antibody reacts specifically with Goat IgG and with light chains common to other Goat immunoglobulins. No antibody was detected against non immunoglobulin serum proteins. Reduced cross-reactivity to chicken, horse, human, mouse, pig, rabbit and rat was detected.
アプリケーション	適用あり: Flow Cyt, IHC-P, ICC/IF, WB
吸着処理血清	Chicken, Horse, Human, Mouse, Pig, Rabbit, Rat more details
標識	DyLight® 488. Ex: 493nm, Em: 518nm

製品の特性


製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C.
バッファー	pH: 6.8 Preservative: 0.09% Sodium azide Constituents: 0.2% BSA, PBS
精製度	Immunogen affinity purified
特記事項 (精製)	Antiserum was cross adsorbed using chicken, human, mouse, pig, rabbit and rat immunosorbents to remove cross reactive antibodies. This antibody was isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to DyLight® 488.
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、**次のテスト済みアプリケーションにおけるab96935の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt		1/50 - 1/200.
IHC-P		1/50 - 1/500.
ICC/IF		1/50 - 1/500.
WB		1/1000 - 1/2000. Predicted molecular weight: 36 kDa. 5% non-fat dry milk in PBST or TBST is recommended for blocking and incubation of antibodies. BSA is not recommended.

画像




All lanes : cocktail of mouse anti-hnRNP
, rabbit anti-actin, and goat anti-GAPDH at 1 µg/ml

All lanes : Hela

Secondary
All lanes : cocktail of DyLight® 680-conjugated donkey anti-mouse
(green), DyLight® 800-
conjugated donkey anti-rabbit (red), and DyLight® 488-conjugated
donkey anti-goat (blue) (ab96935) at 0.5 µg/ml


Predicted band size: 36 kDa



Immunohistochemistry (PFA perfusion fixed frozen sections) - Donkey Anti-Goat IgG H&L (DyLight® 488) preadsorbed (ab96935)

Image is courtesy of an AbReview submitted by Ms Françoise Geffroy.

Immunohistochemical analysis of Paraformaldehyde perfusion-fixed frozen human neuroblastoma xenograft into a murine host. The image is labelling LIMP2 with **ab87637** at 20°C for 1 hour at a dilution of 1/50 with 5% Donkey Serum. Permeabilization was done with 0.2% Triton. Secondary used was ab96935 at a 1/250 dilution.



Donkey Anti-Goat IgG H&L (DyLight® 488) preadsorbed (ab96935)

Emission spectra of DyLight® fluorochromes available in our catalog.

Line colors represent the approximate visible colors of the wavelength maxima.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors